

P-ISSN: 2706-7483
E-ISSN: 2706-7491
IJGGE 2020; 2(2): 23-28
Received: 14-05-2020
Accepted: 27-06-2020

M Reshetnyk
Ph.D., Researcher,
National Museum of Natural
History Nasu, Kiev, Ukraine

Rock magnetic properties of enderbites in Gaisin Block of Ukrainian Shield

M Reshetnyk

Abstract

The actual problems of detail studying of geological structures of the Ukrainian Precambrian Shield (US) were used. The aim of the work was to verify the possibility mapping of xenolites of enderbites of the abnormal magnetic field. The magnetic susceptibility Curie points of ferromagnetic minerals and other properties of enderbites were investigated. It is shown that the studies enderbites not create high positive anomalies of the magnetic field, and are actually "dumb" in the anomalous magnetic field. On the prospect of suggested further study of those areas where high positive magnetic field anomalies coincide with places spread enderbite.

Keywords: rock magnetic properties, enderbites, Ukrainian shield

1. Introduction

In expositions of *Ukrainian Precambrian Shield (US)* due to magnitometric methods during state geological testing in 1960-1980 years were practically not investigated. Now, an actual problem of geological re-carting of US our government is declared [7]. The investigation of the US with new methods based on more detailed and complex geology-geophysical become actual. In this article authors new detailed complex magnitometric methods [method MMS] are used for investigation xenolites of the enderbites.

The basic question of article is how enderbites are sited in Gaisin block of US (along rivers Southern Boug). It is known [31] because rocks in Gaisin block were cardinally changed during metamorphism but enderbites is very ancient rocks that weren't changed. Therefore, studying of enderbites magnetic properties is important for fundamental investigation of Precambrian rocks - that is result of appearance and the first formation of the Earth's crust.

The rocks in the US are mixed in a difficult correlation with each other. It is very difficult to set on map the boundaries of various rocks, because they very rarely come to the surface and are almost invisible. Almost all Precambrian basement of US is hidden and covered by layers of chalk, clay, sandstone.

The geologists-cartographers detected the Precambrian basement through abnormal magnetic field induction and gravitational anomalies analysed. Sediments rocks hasn't many magnetite minerals, therefore their rocks do not produced anomalies of magnetic field. But the different granites of US can have a different numbers of ferromagnetic minerals and formed different intensive magnetic field anomalies. The old methods for interpretation abnormal magnetic field are used different petrographic groups it has some magnetic properties of samples for all area. Today, investigation samples with concrete exposures (different petrographic composition) in complex with observation magnetic field in this territory [complex method] and this material are used for complex geological analysis of abnormal magnetic field.

Unchanged enderbites xenolites in Gaisin block are present conserved. Enderbite consists practically from all granite-forming minerals. There are a quartz, hypersthene, biotite and feldspar. The difference is in content of feldspar – in enderbite he not consist kaly K. This is formed green-gray colour enderbite, whenever granites are gray or pink. Enderbite has unique macrostructures [8, 34].

2. Material and Methods

Study area. In this article investigation area in the US Ros-Tykych megablock in the Uman small blocks in the Gaisin very small blocks. Gaisin blocks between Gayvoron and Podol block (parts of Dnistrovo-Bug megablock). Nemirov and Obodnov fractures Gaisin from Podol block is separated. Dashev fracture Gaisin from all other Uman block is separated.

Corresponding Author:
M Reshetnyk
Ph.D., Researcher,
National Museum of Natural
History Nasu, Kiev, Ukraine

Dashev fracture Gaisyn from all other Uman block is separated (Figure 1).




Fig 1: Geological structure position of investigation area is big square in global geological structure of Ukrainian shield

Gaisyn block consists of unique composite rocks, it is the Gaisyn complex [30, 31]. Gaisyn complex it is very motley because plagiogranites and grano-diorites are alternated in one metre and contains xenolites of gneiss, kristal slate, enderbyite. Xenolites of enderbyite is unique rocks in Gaisyn complex. On all territory of Gaisyn block is one exposure of enderbyites. Enderbyites exposure located on board of river Gorodishe (to north of the village Sytkivci, Vinniza region

of Ukraine, coordinates latitude 48 55 42.01 N, longitude 29 11 30.51 E). It exposure of enderbyites in this article is investigated. The enderbyites exposure located on Sutkivci territory geological mapping. Katuk and all Sutkivci territory were geological investigated and created geological map in scale 1:10000 [4]. We geological map with magnetic anomalies field map are combined (Figure 2).




Fig 2: Fragment of geological map combined with anomalies magnetic field T map of area among villages Vysha Kropyvnia, Rubigne and Sytkivci (according [4]): 1 – rose granites, 2 – enderbyites, 3 – crystals shale, 4,5,6 – isolines of high and low T in nanoteslas nTl, 7 – fracture valid and from geophysical date, 8 – isolines of high gravity field mG, black point – sites of enderbyites exposure

On figure 2 we can see then big xenolites of enderbyites are sited on different anomalies magnetic field where there is high T (up to 1500 nTl) and low T (-500 nTl). Thus, according [4], the source of negative and positive anomalies of magnetic field are same enderbyites. Therefore is actual to

study the magnetic properties of enderbyites to determine the nature of such ambiguity reflected in the abnormal magnetic field.

Methodology. In this work the authors new complexing detailed magnetometric method is using by enderbyites

investigation. It is method of magnetic scanning (MMS) which let use a new possibilities of mapping for high-differentiated sectors of precambrian basement. Magnetic Scanning – it is a complex petro-magnetic method of detail investigation of rocks exposures as we described earlier [19-20]. In this method, laboratoric and “in field” magnetometric investigations are carried out in direct interrelation with each other.

At the first stage authors analyzed and correlations “magnetic field - geological structure” appears.

At the second stage, magnetometric investigations “in field” are carried out. The detailed measuring of magnetic susceptibility MS were executed by capameter KT-5 (Czech production). The modern counterparts of KT-5 is KT-9. The templates of rose granites and enderbytes were selected. The magnetic susceptibility of their templates were measured on astatic laboratory magnetometer LAM-24 (Czech production). The principle of the sensor is a magnetic field of pattern shifts magnets of astatic system from the equilibrium. It is need to measure in 12 direction and how much deviated astatic system. These measurements allow to calculate the value of the natural remanent magnetization of the sample, its magnetic susceptibility and orientation of remanent magnetization vector.

The magnetic minerals in rocks are definite by thermomagnetic analysis. It consists in consequent

measuring of magnetic susceptibility MS of heated template. The laboratoric magnetometer KLY-2 fixes a change MS of sample which heated in an oven. The modern counterparts of KLY-2 is MFK1 Kappabridges of kompani AGICO.

The density of sample was measured by method of hydrostatic gravimetry.

The dynamic of change of magnetic minerals was investigated by rock microscopy.

3. Results and discussion

Over enderbytes exposure on the board of river Gorodishe T = -351 nTl. We MS of enderbytes on exposure were observed. 80% of MS numbers is small $0-12 \cdot 10^{-3}$ u.SI (fig.3). It isn't big for granitoides of Gaisin complexes. (Then we were investigated other granitoid exposures near enderbytes exposure we were indentified pink granites with small MS $<10 \cdot 10^{-3}$ u.SI and migmatites with big MS $>30 \cdot 10^{-3}$ u.SI.) Some big MS in the range $40-50 \cdot 10^{-3}$ u.SI interspersed in the total weight with low values of MS $0-12 \cdot 10^{-3}$ u.SI. It big MS found on the exposure is not regularly (Figure 4). This can be explained by not-homogenous distribution of magnetite that have a several agglomerates in the enderbyte.




Fig 3: Diagram MS (MS range is $\cdot 10^{-3}$ u.SI) distribution measured by the enderbytes exposure



Fig 4: Limited fragment of enderbytes exposition carrier with markers of sites with MS $40-50 \cdot 10^{-3}$ u.SI

MS very high values in the range $70-75 \cdot 10^{-3}$ u.SI obtained in only three measuring points (MS measured in 620 points on the network with 0.2m increments). MS sample

distribution coincided with the distribution of MS measured at the enderbytes exposure. The most samples (43 samples out of 54) has low MS ($0-12 \cdot 10^{-3}$ u.SI) indicating that a small amount of magnetite. At the same time, thermomagnetic analysis shows that their templates have magnetite and pirotine (Figure 5).




Fig 5: Thermomagnetic curves of templates

The curve MS_i/MS_1 falls sharply after heating to $380 \text{ }^\circ\text{C}$ indicating the presence pirotine. Ore microscopy revealed the presence of pirotine grains in enderbytes. Enderbite samples which was identified pyrrhotite have heightened the Q factor in the range 1.5-2 units (normal 0.4-0.6 units).

These samples has back oriented vector of the remanent magnetization (Figure 6). Mostly vector of the remanent magnetization is oriented in the direction of modern magnetic field (for the coordinates our enderbites exposure declination 11 37.68 inclination 64 66.05).




Fig 6: Stereoscopic of the remanent magnetization vector of enderbite. Black circles - straight, hollow circles - reverse orientation of the vector of the remanent magnetization

The low values of the magnetic susceptibility, and hence the induced magnetization are indicated that enderbite will not significantly affect the nature of the anomalous magnetic field. That part enderbites it is inversely oriented remanent magnetization can only weaken the magnetic field. Map of the anomalous magnetic field shows that the enderbites exposure is in the negative conjugate field anomalies. The positive part of the conjugate anomaly is located south-west enderbites exposure (see Figure 2). It is necessary to examine the method of magnetic scanning exposure in the area of positive anomalies of the magnetic field ($T > 1000nTl$) in order to identify the source of that.

Place high positive anomalies of the magnetic field coincides with the place enderbite body position (see Figure 2). Our research has shown that enderbites can not create high positive anomalies of the magnetic field. This means that the anomaly magnetic field may cause or other rocks or there enderbites are located on the magnetic properties different from those examined enderbites.

Enderbite density ranges from 2.67 to 2.77 g/cm^3 . Enderbite density ranges from 2.67 to 2.77 g/cm^3 while the pink granite have a density of 2.6 g/cm^3 . Migmatites in representation of areas in figure 2 it we invistigeted have a density from 2.6 to 2.7 g/cm^3 . This indicates that enderbites have high density and can produce positive anomalies in the gravitational field. Compare anomalies card Bug and geological map shows that the anomalies coincide with the contours of the body contours enderbite. This suggests that the creation of a geological map does not take into account the anomalous magnetic field map. This requires a new analysis of existing spatial information.

4. Conclusion

The magnetic susceptibility of enderbites in Gaisin Block are investigated in the first time. It is established that enderbites are weak-magnetic ($MS < 12 \cdot 10^{-3}$ u.SI) rocks with magnetite+pyrotine ferromagnetic minerals.

Experimental data can show that enderbites cannot produce a high induction of magnetic field. On the prospect of suggested further study of those areas where high positive magnetic field anomalies coincide with places spread enderbite.

5. References

1. Antsiferov AV, Dovbnich MM, Tyapkin KF etc. Integration of geophysical methods, Donetsk: Weber. 2008, 335.
2. Chmielewska VK. Integration of geophysical methods for studying the Earth's crust, International University of Nature, Society and Man "Dubna" (1997), <http://geo.web.ru/db/msg.html?mid=1161636&uri=pag e200011.html>
3. Instruction of magnitometrics, http://users.monash.edu.au/~rjarmit/Mag_sus_meter_in structions/KappaMeter_KT9/KT9_manual.pdf
4. Katyuk *et al.* Group Geological Survey 1: 50,000 with the general search area sheets of M-35-119 A, B - 120, B, (Gysin), Report geol.- shooting squad for 1987-1991. Kiev, 1992, 345.
5. Mondal S, Piper JDA, Hunt L, Bandyopadyay G, Basu Mallik Paleomagnetic S. Rock magnetic study of charnockites from Tamil Nadu, India, and the Ur protocontinent in Early Paleoproterozoic times, Precambrian Research Journal of Asian Earth Sciences, 2009; 34: 493-506.
6. Lysek A *et al.* Formational analysis the lower range of the Precambrian of the Ukrainian Shield during geological survey, Kiev, 2006, 203.
7. Lisak AM, Pascenco VG. Sobitovaya formation is the western part of the Ukrainian shield (volume and internal structure), Questions of the theory and practice of conformational studies of the Precambrian lower, Lviv: Vishcha School, 1981, 92-104.
8. Organization and realization geological reinvestigation mapping area scale of 1: 200,000, complication and preparation to edition state geological map of Ukraine scale of 1: 200,000, Instruction, Kiev, 1999, 296.
9. Kostenko M. Geological studying of early mapped surfaces, and formation of State Geological Map 1:200 000 ..., Mineral.Resoursy of Ukraine, 2018; v.3, 3-12.
10. Manshtein A.K. Small-deep geophysics. Novosibirsk edition (2002), 136 pages. <http://www.nemfis.ru/maloglgf.pdf>
11. Murzviov L. Possibilities of magnetometric scanning in continuous regime at investigation of petrol-perspective sector. Neftegazovoe Delo (Petrol-gaz Business, in Russian), 2007. <http://www.ogbus.ru>.
12. Starokadomsky D, Ishenko A, Reshetnyk M. Filling with the Graphene Nanoplates as a Way to Improve Properties of Epoxy-Composites for Industrial and Geophysical Machinery// American Journal of Physics and Applications – 2017; 5(6):120-125.
13. JDA Piper, Basu Mallik S, Bondyopadhyay G, Mondal S, Das AK. Palaeomagnetic and Rock Magnetic study of a deeply-exposed continental crustal section in the charnockite Belt of southern India: implications to crustal magnetization and palaeoproterozoic continental nuclei, Precambrian Res. 2003; 121:185–219.
14. Reshetnyk MN. The complex magnetic scanning as an effective method to investigate the exposures of Precambrian Basement: example of Ukrainian Shield, Scientific Journal of Pure and Applied Sciences. 2012; 1(1):22–29. <http://www.sjournals.com/index.php/SJPAS/article/vie w/239/pdf>
15. Reshetnyk MM, Sukhorada AV, Khomenko RV. Structural and geological information of magnetic

- scanning of the Precambrian foundation (at the Middle Shield of the Ukrainian Shield, in the region of Gayvoron-Zavalla), News of the Kiev National University of Taras Shevchenko. Geology, 2010; 48:44–48.
16. Reshetnyk MN. Scanning of Magnetic Properties in Precambrian Basement, Geosciences, 2012; 2(3):33–38. <http://article.sapub.org/10.5923.j.geo.20120203.01.html>
 17. MM Reshetnyk Rocks magnetic properties of Precambrian basement (for example Ukrainian shield) structural-geological informativity. Diss. PhD speciality geophysics, T.Shevchenko National University of Kyiv, Kiev, 2012, 200.
 18. Reshetnyk MM. Small-scale magnetometry at the Polish geological structures of the Precambrian foundation, News of the Kiev National University of Taras Shevchenko. Geology, 2013; 1(60):51-55.
 19. Zavovskiy V. Mountain-sun compass. Geophysical Journal. 2003; 25(5):112–117.
 20. Ryabokon VV, Scherbakov IB. Sobit Ukrainian shield and their genesis, Kiev, Preprint IGFM, 1977, 53.
 21. Sherbakov I. Petrology of Ukrainian Shield, Lvov, ZUKZ, 2005, 362.
 22. Sukhorada AV, Reshetnik MM, Khomenko RV. Magnitni markers structuring kristalichnogo foundation Serednogo Pobuzhzhya (on prikladi area Haivoron-Zavallia), Visn. Kiev. the University. Ser. "Geologiya", 2008; 44:17-20.
 23. Sukhorada AV, Guziy MI, Ivahnenko OP, Popov SA. Before boarding about magnitomineralogichnu model Gaysinskogo block Ukrayinsky shield, News of the Kiev National University of Taras Shevchenko. Geology, 1995; 14:103-115.
 24. Zhang J, JDA. Piper Magnetic fabric and post-orogenic uplift and cooling magnetisations in a Precambrian granulite terrain: The Datong-Huai'an region of the North China Shield, *Tectonophysics*, 1994; 234: 227–246.
 25. Kujelow TK. Geological structure of Precambrian in Middle Pobujie region. Geological Journal (USSR, in Russian), 1960, 20(2): 30–46.
 26. Kujelow TK. Reflection of the tectonics of the Ukrainian Shield in magnetic and gravitational fields, Soviet Geology. 1964; 2:75–87.
 27. Orlyuk MI, Orlova MI. A review of the results of studies of the paleo-magnetic information content of Precambrian crystalline rocks of the Ukrainian Shield. Geophysical Journal. 2013, 84-108.
 28. Tyapkin KF. Study of fault and folded structures of the Precambrian by geological and geophysical methods. - Kiev: Naukova Dumka Edition, 1986, 168.
 29. Tyapkin KF, Dovbnich MM. A new rotational hypothesis of structure formation and its geological and mathematical substantiation: monograph. Donetsk: Knowledge Edition, 2009, 342.
 30. Tolstoy MI. Petrophysics of granitoids of the Ukrainian shield [MI Tolstoy, AV Chekunov, IB Shcherbakov]. Kiev: Naukova Dumka Edition, 1987, 232.
 31. Fedyuk V.I. Micromagnetic survey and its geological capabilities. Moscow: Gostekhizdat Edition, 1958, 63.
 32. Sholpo LE. The use of magnetism of rocks in geological survey [Sholpo LE, Rusinov B.Sh., Ilaev M.G. and etc.]. Leningrad: Nedra Edition, 1986, 234.
 33. Kosnyreva MV. Development of a complex of geophysical methods for solving applied problems of soil mapping: author. dis. on the science. Degree of Ph.D. geol.-miner.: spec. 04.00.22 "geophysicist". Moscow, 2007, 201.
 34. Starokadomsky D, Ishenko A, Reshetnyk M. Filling with the Graphene Nanoplates as a Way to Improve Properties of Epoxy-Composites for Industrial and Geophysical Machinery. American Journal of Physics and Applications 2017, 5(6):120-125.
 35. Starokadomsky D, Reshetnyk M. Microfilled Epoxy-composites, capable of thermo-hardening and thermo-plasticization after hard heating (200-300 oC) - for "in-field/offroad" use in bio-,agro-, medservice. Biomedical J.of Scientific & Technical Research (ImpactF=0,548) 19(1): 14118-14123 – <https://biomedres.us/submit-manuscript.php>
 36. Завойський В.М. Гірничий сонячний компас // Геофізический журнал. – 2003; 25(5):112–117.
 37. Ryabokon VV, Scherbakov Sobit IB. Ukrainian shield and their genesis, Kiev, Preprint IGFM, 1977, 53.
 38. Sherbakov I. Petrology of Ukrainian Shield, Lvov, ZUKZ, 2005, 362.
 39. Sukhorada AV, Reshetnik MM, Khomenko RV. Magnitni markers structuring kristalichnogo foundation Serednogo Pobuzhzhya (on prikladi area Haivoron-Zavallia), Visn. Kiev. the University. Ser. "Geologiya", 2008; 44:17-20.
 40. Sukhorada AV, Guziy MI, Ivahnenko OP, Popov SA. Before boarding about magnitomineralogichnu model Gaysinskogo block Ukrayinsky shield, *Visn. Kiev. the University. Ser. "Geologiya"*, 1995; 14:103-115.
 41. Zhang J, J.D.A. Piper Magnetic fabric and post-orogenic uplift and cooling magnetisations in a Precambrian granulite terrain: The Datong-Huai'an region of the North China Shield, *Tectonophysics*, 1994; 234: 227–246.
 42. Кужелов ТК. Геологічна структура докембрію Середнього Побужжя // Геологічний журнал. – 1960; 20(2):30–46.
 43. Кужелов Т.К. Отражение складчатой тектоники Украинского щита в магнитном и гравитационном полях// Советская геология. – 1964, 75–87.
 44. Орлюк М.И., М.И.Орлова. Обзор результатов исследований палео-магнитной информативности докембрийских кристаллических пород Украинского Щита. Геофізический Журнал, 2013, 84-108.
 45. Тяпкин К. Ф. Изучение разломных и складчатых структур докембрия геолого-геофізическими методами. – Киев : Наук. думка, 1986, 168.
 46. Тяпкин К.Ф., Довбнич М.М. Новая ротационная гипотеза структурообразования и ее геолого-математическое обоснование: монография. – Донецк: «Ноулидж», 2009, 342.
 47. Толстой М.И. Петрофизика гранитоидов Украинского щита [М.И.Толстой, А.В.Чекунов, И.Б.Щербаков]. – К., 1987, 232с.
 48. Федюк В.И. Микромагнитная съемка и ее геологические возможности. – М: – Гостехиздат, 1958, 63с.
 49. Шолпо Л.Е. Использование магнетизма горных пород при геологической съемке [Шолпо Л.Е., Русинов Б.Ш., Илаев М.Г. и др.]. – Ленинград. : Недра, 1986, 234с.

50. Коснырева МВ. Разработка комплекса геофизических методов для решения прикладных задач почвенного картирования : автореф. дис. на здобуття наук. степеня канд. геол.-мін.: спец. 04.00.22 «геофізик» / М.В. Коснірева. – М, 2007, 20[1] с.