

P-ISSN: 2706-7483  
E-ISSN: 2706-7491  
IJGGE 2022; 4(1): 84-88  
Received: 02-01-2022  
Accepted: 05-02-2022

**Mushtaq Ahmad Ganie**  
Department of Environmental  
Science and Limnology,  
Barkatullah University,  
Bhopal, Madhya Pradesh,  
India

**Hilal Ahmad Bhat**  
Department of Environmental  
Science and Limnology,  
Barkatullah University,  
Bhopal, Madhya Pradesh,  
India

**Abhilasha Bhawsar**  
Department of Environmental  
Science and Limnology,  
Barkatullah University,  
Bhopal, Madhya Pradesh,  
India

**Corresponding Author:**  
**Mushtaq Ahmad Ganie**  
Department of Environmental  
Science and Limnology,  
Barkatullah University,  
Bhopal, Madhya Pradesh,  
India

## Evaluation of anthropogenic stress on ecological quality of riparian habitat of Kolar River

**Mushtaq Ahmad Ganie, Hilal Ahmad Bhat and Abhilasha Bhawsar**

### Abstract

Riparian zones are among the major ecologically valuable ecosystems due to their abundant richness in biological diversity and services. The present study was conducted on Kolar River, in a stretch of about 80 kms to assess the ecological quality of riparian habitat using *qualitat del bosc de ribera* (QBR) index. In this study, QBR index has been used to determine the riparian habitat quality of Kolar River. The selected stretch (60 km) has been divided into eight different sampling sites. The on-site assessment has been carried out on the left and right banks of Kolar River on these sites using different parameters for assessment of riparian habitat quality like total riparian cover, cover structure, cover quality and channel alteration. These parameters have been analyzed and it was evaluated that some sites along the banks of river viz; (S1, QBR score = 75), (S3, QBR score = 85) (S3, QBR score = 80) and shows QBR score in between 75-90 (Green) which reveals good quality and some disturbance of riparian habitat whereas the sites S4, S5, S6, and S8 reflects QBR score of 70, 65, 60 and 55 respectively which lies in between 55-70 (Yellow) that determines fair quality and important disturbance of riparian habitat. Moreover, site seven (S7) shows (QBR score=50) in the range of 30-50 (Orange) which reflects strong alteration and poor quality of riparian zone. The current conditions of riparian zone were due to increase in anthropogenic activities like built-up, overgrazing, deforestation and expansion in agricultural activities. Human dependence as livelihood is being the major cause of the change in riparian zone of study area. Therefore, it is compulsory to save riparian habitat quality of Kolar River from further disturbances and degradation by anthropogenic pressures.

**Keywords:** Riparian habitat, ecosystem, Kolar River, QBR index, anthropogenic

### 1. Introduction

Riparian habitats are important ecological areas having rich biodiversity and provides important ecosystem services (Gurnell *et al.* 2012; DeVore *et al.* 2016, Pandey *et al.* 2022) [25, 26]. The ecologically functional riparian zone (ERZ) supports main ecological functions (Lind *et al.* 2019; Martin and Chambers, 2002; Naiman *et al.* 1993) [27, 28, 12]. The word 'riparian' is derived from Latin word '*riparius*' which means on the bank of a river (Naiman and Decamps, 1997) [29] "A riparian area can be defined as the transitional area between a river or stream and the adjoining upland vegetation, including both the stream channel itself and its surrounding land which influences due to fluctuating water levels" (Corbacho *et al.* 2003; Goebel *et al.* 2003; Seibert and McGlynn, 2005) [14, 16, 30] but they vary from upland in term of their hydrologic and soil characteristics. Riparian zone provides ecosystem services for riverine ecosystem and it also acting an important role for maintaining abiotic and biotic components. It provides shelter and food for fauna living nearby the river and helps in water filtration (Bunn and Arthington, 2002) [31] and aquifer recharge. Riparian zone helps in maintaining water quality, control sediment erosion (Noe & Hupp, 2009, flooding (Cadotte *et al.* 2011) [32, 53] & temperature control, decreasing hydrological risk and construct stable river banks (Fu *et al.* 2017) [15]. Two features that distinguish riparian ecosystems are the hydrologic interaction that occurs between the stream channel and adjacent areas through the periodic exchange of surface and ground water, and the unique geomorphic characteristics and vegetation communities living in response to this hydrologic interaction (Richards 1982; Kovalchik and Chitwood 1990; Gregory *et al.* 1991; Goodwin *et al.* 1997; Malanson 1993) [2, 4, 5, 6, 7].

Riparian areas are considered as one of the most productive ecosystems (Jansson *et al.* 2000) in the world, but are also among the most threatened (Brunotte *et al.* 2009; Dixon *et al.* 2016; Strokal *et al.* 2016; Ltifi *et al.* 2017; Wright *et al.* 2017; Knouft and Ficklin, 2017; Capon *et al.* 2013; Mohit and Samant 2012) [34, 36, 38, 37, 35, 8, 13].


Pressures on rivers and riparian zones are expected to increase in the coming decades due to human population growth, land use change, and climate change (Seager *et al.* 2013) [40] heightening the importance of strategic riparian management (Capon *et al.* 2013, Mello *et al.* 2017) [8, 10]. The reduction and alteration of riparian forests have resulted in habitat fragmentation and destruction of habitat of diversity in the river system (Ding *et al.* 2013; Ou *et al.* 2016; Yang *et al.* 2016; Zhang *et al.* 2013; Kuglerover *et al.* 2014) [43, 39, 41, 42, 44].

Assessment of riparian quality has been globally evaluated from decades by applying different protocols and indices. In this study, QBR index was suitable for the present study area. The QBR index (“*qualitat del bosc de ribera*” or riparian forest quality) is convenient to use field method for evaluation of the habitat quality of riparian forests. This index has been first developed for use in Mediterranean streams in Spain (Munne *et al.* 2003) [17]. This index is based on score and has been divided into four main aspects of the

riparian zone *viz*; total riparian cover, cover structure, cover quality, and channel alteration. Its main focus is to contrast sites, to compare sites, to ideal conditions and to assess the success of restoration of riparian zone/riparian forests (Lakhera *et al.* 2020) [54].

**1.1 Study Area**

The present study has been carried out on Kolar River of Madhya Pradesh (M.P) of India. Kolar River rises in the Vindhya Range of district Sehore and then flows in a south-westerly direction to meet the Narmada River near Nasrullahganj which lies in the Raisen district of M.P. Its total drainage area of Kolar River is of 1,347 km<sup>2</sup> and has spread across the two districts in M.P. The upper part of the river basin lies at an elevation of 350-600 meters from MSL and much of it is under tropical deciduous forest area. The river debouches into the plains near Jholiapur. The map showing study area is given (Fig. 1).



**Fig 1:** Showing location of sampling sites

**2. Material and Methods**

The qualitative assessment has been adopted for the analysis of study area. In this context, the study area has been divided into 8 different sampling sites *viz*; Near Reservoir = S1, Near Filtration Plant = S2, new bridge upper side = S3, new bridge lower side =S4, Chilkapani = S5, Jabra Malkhar = S6, Amdoh = S7 and Near Jholiapur Bridge = S8 (Fig. 1). In the present investigation, the riparian habitat quality of Kolar River was assessed by using QBR Index Munne *et al.* 2003 [17] (Table 1). The QBR (“*qualitat del bosc de ribera*” or riparian forest quality) is convenient to apply in field method for assessing ecological quality of riparian habitat. This protocol was developed by F.E.M. (Freshwater Ecology and Management) research group of *Universitatde Barcelona* (Munne *et al.* 2003) [17]. The QBR index is based on four classes of a riparian habitat *viz.*, total riparian cover, cover structure, cover quality and channel alterations. Each category includes several attributes. The scores for each category range from 0 to 25, being 100 the maximum total

score assigned to the highest quality and, representing five classes (Table 1).

**Table 1:** Quality ranges of riparian habitat as per the QBR Protocol (Munne *et al.* 2003) [17]

Riparian Habitat Quality Class	QBR Score	Colour
Riparian habitat in natural condition	>=95	Blue
Some disturbance, good quality	75-90	Green
Disturbance important, fair quality	55-70	Yellow
Strong alteration, poor quality	30-50	Orange
Extreme degradation, bad quality	<=25	Red


**3. Result and Discussion**

In the present study, riparian habitat quality of Kolar River was assessed by using QBR index, the parameters like total riparian cover, cover structure, cover quality and channel alteration were observed at all the mentioned sites of Kolar River. The study found well differences in the results from one location to another as shown in the table 2 and figure 2.

**Table 2:** Score of Riparian Habitat Quality of Kolar River by Using QBR Index


Categories	Score							
	S1	S2	S3	S4	S5	S6	S7	S8
Total riparian cover	25	20	25	25	20	15	15	15
Cover structure	15	20	20	20	15	15	10	15
Cover quality	15	15	15	15	15	10	10	10
Channel alteration	20	15	25	20	15	20	15	15
Final score	75	70	85	80	65	60	50	55
Riparian Zone Condition and Quality	Some disturbance, good quality	Disturbance important, fair quality	Some disturbance, good quality	Some disturbance, good quality	Disturbance important, fair quality	Disturbance important, fair quality	Strong alteration, poor quality	Disturbance important, fair quality

The analysis of riparian zone of Kolar River has been carried out on eight sampling sites using QBR protocol.



**Fig 2:** QBR score of riparian habitat quality of selected sites

According to the protocol (QBR index), strong alteration and poor quality of riparian zone was observed on one site (S7), little disturbance and fair quality of riparian zone was found in four sites (S2, S5, S6 & S8) with important disturbances and three sites (S1, S3, & S4) were found in good quality and some disturbances were observed in these areas (Figure 3).




**Fig 3:** Final QBR score of riparian habitats at different sampling sites of Kolar River

On inclusive observation of the riparian zone, it has been found that 44% area of riparian zone was of good quality and possess some disturbance whereas, 46% area of riparian zone fall under little disturbance and of fair quality and remaining 10% shows poor quality and strongly altered riparian areas (Fig. 4). According to the study, disturbances on the riparian vegetation occur due to the changes in the land use pattern. Agricultural activities, overgrazing and

deforestation are the main activities observed in the riparian area. The cultivation of crops on the banks of river is responsible for degradation of riparian area. Identical observations were reported in river Narmada using QBR index for assessing riparian zone quality on 25 stations in which, 13 stations fall under bad quality, 7 stations were found in fair quality, 4 stations showed very bad quality while only one station was in good quality of riparian zone (Kumar *et al.* 2019) [46]. Tembhare *et al.* (2018) [21] assessed the ecological quality of riparian habitat of streams of Narmada River basin using QBR index to evaluate the riparian habitat quality of Barna and Jamner streams during the carried-out study on the left and right banks of the streams. They observed that Barna stream showed fair riparian quality due to the presence of a large number of trees whilst, Jamner stream fall under bad riparian quality due to anthropogenic pressure.

The fifth longest river of China, Songhua River, which showed 60% of the riparian zone, was disturbed by human activities like build-up and farm land contraction (Bolin *et al.* 2017) [47]. Degradation of riparian zone in Kaliyadeh stream the tributary of River Narmada in the central zone was reported by Pandey *et al.* (2015) [48], the same was observed at Bhahner River by Bahsir *et al.*, (2015) [49] and Chandni Nalla by Chaurasia, *et al.*, (2015) [50]. Expansion of agriculture practices on the banks of Chandni Nalla was also reported by War *et al.*, (2014) [51] which are more responsible for soil erosion and ecological degradation of a stream/River. Poor qualities of riparian zone of River Narmada through the dominance of agricultural activities were also stated by Vyas *et al.*, (2012) [52].



**Fig 4:** Graphical representation of status of riparian quality

#### 4. Conclusion

In this study the riparian habitat quality assessment has been evaluated using QBR index for assessing the different parameters such as total riparian cover, cover structure, cover quality and channel alteration. It was observed that some of the selected locations represent good quality, some locations reflect fair and some signifies poor riparian habitat quality. It has been suggested that the riparian vegetation cover of Kolar River needs improvement and rejuvenation besides plantation. Moreover, this study clearly predicts that the land use pattern and riparian conditions have profound effect in determining riparian habitat quality. So, it becomes compulsory to protect the riparian habitat quality of Kolar River from further disturbances and degradation by anthropogenic pressures. Therefore, government should categorise this area as protected area and give necessary attention for restoration of aesthetic value and well planning for maintain overall quality of Kolar River. Moreover, proper management and policies should be framed for the settlement of human habitation along the buffer zones of the river network which will help in saving the ecology as well as aesthetic value of Kolar River.

#### 5. References

- Williams GP. Bank full discharge of rivers. *Water Resources Research*. 1978;(14):1141-1154.
- Richards K. *Rivers: Form and Process in Alluvial Channels*. Methuen and Company, London, England, 1982, 358.
- Harris RE. Occurrence of vegetation on geomorphic surfaces in the active floodplain of a California alluvial stream. *American Midland Naturalist*. 1987;(118):393-405.
- Kovalchik BL, Chitwood LA. Use of Geomorphology in the Classification of Riparian Plant Associations in Mountainous Landscapes of Central Oregon, USA. *Forest, Ecology and Management*. 1990;(33-4):405-18.
- Gregory SV, Swanson FJ, McKee WA, Cummins KW. An ecosystem perspective of riparian zones. *Bioscience*. 1991;(41):540-551.
- Goodwin CN, Hawkins CP, Kershner JL. Riparian Restoration in the Western United States: Overview and Perspective. *Restoration Ecology*. 1997;(5):4-14.
- Malanson GP. *Riparian Landscapes*. Cambridge University Press, New York, NY, 1993.
- Capon SJ, Chambers LE, Nally Mac R, Naiman RJ, Davies P, Marshall N, *et al*. Riparian ecosystems in the 21st century: hotspots for climate change. *Ecosystems*. 2013;(16):359-381.
- Martinez-Fernandez V, Oorschot M, Van, Smit J, De Gonzalez M, Buijse AD. Modelling feedbacks between geomorphological and riparian vegetation responses under climate change in a Mediterranean context. *Earth Surf. Process. Landforms*. 2018;(43):1825-1835.
- Mello K, De Randhir TO, Aversa R, Alberto C. Riparian restoration for protecting water quality in tropical agricultural watersheds. *Ecol. Eng.* 2017;(108):514-524.
- Tockner K, Stanford JA. Riverine flood plains: present state and future trends. *Environ. Conserv.* 2002;(29):308-330.
- Naiman RJ, Decamps H. The ecology of interfaces: riparian zones. *Annual Review of Ecology Systematics*. 1997;(28):621-658.
- Mohit SA, Samant JS. Impact of land use changes on riparian habitats in Panchganga river system. *Proceedings of International Conference, SWDM*, 2012, 203-207.
- Corbacho C, Sanchez JM, Costillo E. Patterns of structural complexity and human disturbance of riparian vegetation in agricultural landscapes of a Mediterranean area. *Agriculture, Ecosystems and Environment*. 2003;(95):495-507.
- Fu B, Li Y, Wang Y, Campbell A, Zhang B, Yin S, *et al*. Evaluation of riparian condition of Songhua River by integration of remote sensing and field measurements. *Scientific Reports*. 2017;7:2565.
- Goebel PC, Hix DM, Semko-Duncan ME. Identifying reference conditions for riparian areas of Ohio. *Special Circular 192*. Wooster, OH: Ohio Agricultural Research and Development Centre, 2003.
- Munne A, Prat N, Sola C, Bonada N, Rieradevall M. A simple field method for assessing the ecological quality of riparian habitat in rivers and streams: QBR index. *Aquatic Conservation: marine and Freshwater Ecosystems*. 2003;(13):147-163.
- Tembhare DA, Renear RB, Mostaghimi S, Lee D. Vegetative filter strips for agricultural nonpoint source pollution control. *Transactions of the American Society of Agricultural Engineers*. 1989;(32):513-519.
- Quadar J, Bhat MA, Vyas V, Shrivastav P, Bhawsar A. Physical habitat assessment of two pristine and polluted streams of Narmada River basin. *Journal of Pharmacy and Biological Science*. 2017;12(6):60-63.
- Barbour MT, Gerritsen J, Snyder BD, Stribling JB. *Rapid bioassessment protocols for in streams and wadeable rivers*. United States Environmental protection agency, Washington, D.C., 1999.
- Tembhare R, Bhawsar A, Vyas V, Bhat MA. Assessing ecological quality of riparian habitat of streams of Narmada River basin using QBR index. *International Journal of Applied and Natural Sciences*. 2018;7(2):87-90.
- Stefunkova Z, Neruda M, Vasekova B. Impact Evaluation of Riparian Vegetation on Aquatic Habitat Quality of Rivers. *Earth and Environmental Science*, 2019, 221.
- Olokeogun OS, Ayanlade A, Popoola OO. Assessment of riparian zone dynamics and its flood-related implications in Eleyele area of Ibadan, Nigeria. *Environ. Syst. Res*. 2020;9:6.
- Corenblit D, Baas A, Balke T, Bouma T, Fromard F,

- Garofano-Gomez V, *et al.* Engineer pioneer plants respond to and affect geomorphic constraints similarly along water–terrestrial interfaces worldwide. *Glob. Ecol. Biogeogr.* 2015;24:1363-1376. DOI: 10.1111/geb.12373.
25. DeVore RM. Elk resource selection patterns in a semiarid riparian corridor. *J Wildl. Manag.* 2016;80(3):479-489. DOI: 10.1002/jwmg.1040.
  26. Pandey S, Kumari T, Verma P, Singh R, Raghubanshi AS. Impact of anthropogenic stresses on riparian ecosystem and their management perspectives. In *Ecological Significance of River Ecosystems*. Elsevier, 2022, 299-324.
  27. Lind L, Hasselquist EM, Laudon H. Towards ecologically functional riparian zones: A meta-analysis to develop guidelines for protecting ecosystem functions and biodiversity in agricultural landscapes. *J Environ. Manage.* 2019;249:109391.
  28. Martin D, Chambers J. Restoration of riparian meadows degraded by livestock grazing: above and belowground responses. *Plant Ecol.* 2002;163:77-91.
  29. Naiman RJ, Decamps H, Pollock M. The role of riparian corridors in maintaining regional biodiversity. *Ecol. Appl.* 1993;3:209-212.
  30. Seibert J, McGlynn BL. Landscape Element Contributions to Storm Runoff. *Encyclopaedia of Hydrological Sciences*. Wiley, Hoboken, 2005.
  31. Bunn SE, Arthington AH. Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environ. Manage.* 2002;30:492-507.
  32. Noe GB, Hupp CR. Retention of riverine sediment and nutrient loads by coastal plain floodplains. *Ecosystems.* 2009;12:728-746.
  33. Jansson R, Nilsson C, Dynesius M, Andersson E. Effects of river regulation on river-margin vegetation: a comparison of eight boreal rivers. *Ecol. Appl.* 2000;10(1):203-224.
  34. Dixon SJ, Sear DA, Odoni NA, Sykes T, Lane SN. The effects of river restoration on catchment scale flood risk and flood hydrology. *Earth Surface Proc Landforms.* 2016;41(7):997-1008.
  35. Knouft JH, Ficklin DL. The potential impacts of climate change on biodiversity in flowing freshwater systems. *Ann. Rev. Ecol. Evol. System.* 2017;48:111-133.
  36. Stokal M, Ma L, Bai Z, Luan S, Kroeze C, Oenema O, *et al.* Alarming nutrient pollution of Chinese rivers as a result of agricultural transitions. *Environ. Res. Lett.* 2016;11:024014.
  37. Wright IA, Belmer N, Davies PJ. Coal mine water pollution and ecological impairment of one of Australia's most 'protected' high conservation-value rivers. *Water Air Soil Pollution.* 2017;228:90.
  38. Ltifi A, Saidi N, Rabhi M, Sealey KS, Hafiane A, Smaoui A, *et al.* Spatial and Temporal variation of parameters in Wadi Andlou, Tunisia-Pollution by pulp mill discharge. *Clean-Soil, Air, Water.* 2017;45(10):1500471.
  39. Ou Y, Wang X, Wang L, Rousseau AN. Landscape influences on water quality in a riparian buffer zone of drinking water source area, Northern China. *Environ. Earth Sci.* 2016;75(114):1-13.
  40. Seager R, Ting M, Li C, Naik N, Cook B, Nakamura J, *et al.* Projections of declining surface-water availability for the southwestern United States. *Nat. Climate Chang.* 2013;3(5):482-486.
  41. Yang H, Wang G, Wang L, Zheng B. Impact of land use changes on water quality in headwaters of the Three Gorges Reservoir. *Environ. Sci. Pollut. Res.* 2016;23:11448-11460.
  42. Zhang Y, Xia J, Shao Q, Zhai X. Water quantity and quality simulation by improved SWAT in highly regulated Huai River Basin of China. *Stoch. Env. Res. Risk A.* 2013;27:11-27.
  43. Ding S, Zhang Y, Liu B, Kong W, Meng W. Effects of riparian land use on water quality and fish communities in the headwater stream of the Taizi River in China. *Front. Environ. Sci.* 2013;7(5):699-708.
  44. Kuglerová L, Agren A, Jansson R, Laudon H. Towards optimizing riparian buffer zones: ecological and biogeochemical implications for forest management. *Forest Ecol. Manag.* 2014;334:74-84.
  45. Pascacio ED, Argueta AO, Castillo-Uzcanga MM, Marcial MR. *Botanical Sciences.* 2018;96(2):180-199. DOI: 10.17129/botsci.1858.
  46. Kumar A, Lakhera K, Vyas V. *International Journal of advance forest science and management.* 2019;3(1):55-60.
  47. Bolin FU, Ying LI, Yeqiao Wang, Anthony Campbell, Bai Zhang, Shubai Yin, *et al.* Evaluation of Riparian Condition of Songhua River by integration of remote sensing and Field measurements. *Scientific Reports.* 2017;7:2565. DOI:10.1038/s41598-017-02772-3.
  48. Pandey S, Kumar A, Vyas V. *Science Insights: An International Journal.* 2015;5(1):1-8. ISSN: 2277-3835.
  49. Bahsir T, Kumar A, Vyas V. *International Journal of Research in Natural and Applied Science.* 2015;5(1):1-6.
  50. Chaurasia A, Kumar A, Vyas V. *International Journal of Research in Environmental Science and Technology.* 2015;5(1):1-9. ISSN: 2249-9695.
  51. War SA, Kumar A, Vyas V. *Advances in Applied Science Research.* 2014;5(2):102-110. ISSN:0976-8610.
  52. Vyas V, Kumar A, Wani SG, Parashar V. *International Journal of Environmental Science.* 2012;3(1):659-674. ISSN:0976-4402. DOI: 10.6088/ijes.2012030131064.
  53. Cadotte MW, Carscadden K, Mirotchnick N. Beyond species: functional diversity and the maintenance of ecological processes and services. *J Appl. Ecol.* 2011;48:1079-1087. DOI: 10.1111/j.1365-2664.2011.02048.x.
  54. Lakhera K, Gurjar P, Vyas V. Ecological Status of Riparian Zone of Sip River: A Tributary of River Narmada in Central Region. *Indian J Applied & Pure Bio.* 2020;35(1):25-31.