

P-ISSN: 2706-7483
E-ISSN: 2706-7491
IJGGE 2023; 5(1): 32-43
Received: 03-11-2022
Accepted: 09-12-2022

Mamta Yadav
India Meteorological
Department, Bhopal, Madhya
Pradesh, India

Vineesha Singh
Department of Earth Sciences,
Barkatullah University
Bhopal, Madhya Pradesh,
India

Corresponding Author:
Mamta Yadav
India Meteorological
Department, Bhopal, Madhya
Pradesh, India

Long-term climatic trends in rainfall pattern analysis over Madhya Pradesh

Mamta Yadav and Vineesha Singh

DOI: <https://doi.org/10.22271/27067483.2023.v5.i1a.139>

Abstract

Climate change is a long-term process and main issue of research now a days. The rainfall, its trends and variability are also affected by climate change and hence the study of rainfall, variability and future predictions are very much important for planning and management.

MP is the second largest state, in India and the fifth largest state in population. Its total area is 308,252 km² and occupies 9.38% area of the country comprises of two meteorological subdivisions viz. East MP & West MP and ten agro-meteorological zones. The varying climatic conditions along with various geographical features have been responsible for drought and flood conditions in different parts of the state. Increasing population and urbanization of the state are causing adverse impacts on the natural environment. The long-term trends of rainfall are required for the planning of optimum adaptation strategies for sustainable use of water. The rainfall trends, its changes and variability are very important for understanding the impact of climate change at regional scale. The study focuses on temporal variation in rainfall at annual, seasonal & monthly scale over East & West MP meteorological subdivisions and MP state as a whole during the period 1871-2016. The long-term changes in rainfall are examined by Mann Kendall and Sen's slope estimates and also analysed the nature relationship with Southern Oscillation Index (SOI) and North Atlantic Ocean (NAO) index. In monthly trends the rainfall over West MP has shown a significant decrease during June while a significant increase during August, whereas the seasonal and annual trends of rainfall are not significant in West MP. In East MP, annual as well as seasonal rainfall during monsoon season have shown significantly decreasing trend which is most prominent in June and July months. Monsoon rainfall is showing significant decreasing trend at 0.1 level of significance along with significant decrease in June month at 0.05 level over the MP state as a whole. The winter and pre-monsoon rainfall showed almost constant (non-significant) trends for the East as well as West MP and for MP state as a whole; however the post monsoon rainfall showed a significantly decreasing trend over West MP at 0.1 level and over all MP state as a whole at 0.05 level of significance during past 50 years. The state receives major rainfall during monsoon season only and rests of the season contribute only 10% to its annual average rainfall. The decreasing tendency in monsoon rainfall and hence annual rainfall, the urban and semi urban areas of the state are facing problems of water scarcity. The changing rainfall trends during monsoon season are affecting kharif crops productivity. The significantly decreasing monsoon and post monsoon rainfall pattern are causing more dependency of farmers on ground water resources.

Keywords: Trends, Mann-Kendall Test, Sen's slope

Introduction

Rainfall, its variability, long-term trends and prediction of future trends are very crucial for economic and natural resource planning. The location of area with respect to equator and its natural relief features characterize the rainfall and runoff patterns. MP is located in Central India. It is one of the economically fastest growing states in our country. At current prices, the Gross State Domestic Product (GSDP) of MP for 2017-18 was Rs 7, 07,047 cores. The compound annual growth rate is 14.0%. Agriculture is the main source of livelihood for the people of the MP. According to census 2011, 69.8% of total worker (urban and rural) and 85.6% of the total worker in rural areas of MP are dependent on agriculture for their livelihood in the state as compared to 72.3% in rural India. In 2017-18, the state produced 42.14 million metric tons of food grains. The state is one of the major cotton producing states of India. MP is being honored "Krishi Karman Award" for the fifth consecutive year from 2011-12 to 2015-16. Economic survey report of MP (2017-18), state has achieved 18% agriculture growth during the last five years and mainly dependent on rainfall for its water

requirements, range from 60–120 cm in different regions of the state. The state is currently exploiting 46% of the available ground water. Among all 51 districts, ground water availability is under semi-critical 23 districts, critical in 7 districts and over-exploited in eight districts. About 99% of the drinking water needs are being fulfilled with ground water and 90% of the ground water is being used for irrigation purposes. The climate of state is predominately moist sub humid to dry sub humid, semi-arid to dry sub-humid and semi-arid in east, west and central plateau and hills respectively, according to agro-climatic regions of India. Sharma *et al.* (2015) ^[23], the state of agriculture report of MP, annual rainfall received in the state varies from 800 mm in the northern and western regions to 1200 mm in the eastern districts. In some year's rainfall goes much below to the normal. Most of the rainfall is received in the Monsoon season from June to September and about 10% of the rainfall is received in the remaining part of the year. Favourable soil and climatic conditions helped the state to be a leading producer of coarse cereals, wheat, soybean, gram, sugarcane, rice, maize, cotton, rapeseed, mustard, arhar, pulses, oil seeds in the country. Rainfall is an important meteorological parameter, which has direct application to agricultural production and water resources. Various attempts have been made since last century to study seasonal and annual trends in rainfall of the country, central India and MP state. Long-term changes in the Indian monsoon rainfall on regional and local scales have important social and economic consequences Walker (1910) ^[27] did not found any trend in monsoon rainfall during 1841-1908 periods. Kumar *et al.* (1992) ^[35] done a study on spatial and sub-seasonal patterns of long-term trends of Indian summer monsoon rainfall. The monthly and seasonal rainfall was quantified by linear trends. West coast, North Andhra Pradesh and North West India are the areas where monsoon season rainfall trends were increasing whereas East MP and adjoining areas, North East India and some parts of Gujrat the trends were decreasing. August rainfall contribution is more prominent to monsoon seasonal rainfall mainly over west coast and central India. Naidu *et al.* (1999) ^[16] studied annual rainfall trends for (1871-1994) period, the negative trends are seen over west central India, central north India and north-eastern parts of India. The positive trend observed over north-west India, covering Haryana, Punjab, East Rajasthan, West Rajasthan and West MP, an isolated area in the east and the peninsula. Statistical tests show that at the 90% significance level, the negative trend is significant over East MP where the rainfall decreases by 1.09 mm/year. Rajeevan and Guhathakurta (2007) ^[8] analysed the trends and observed no significant trend for monsoon and monsoon months for the country as a whole with possible large variations on the regional scale. The July rainfall has decreased for most parts of the central and peninsular India but increased significantly in the north-eastern parts of the country. August rainfall has increased significantly (at 95% significance level) for the subdivisions Konkan and Goa, Marathwada, Madhya Maharashtra subdivision, Vidarbha, West MP, Telangana and west Uttar Pradesh Rainfall for the subdivisions viz. east Uttar Pradesh, Bihar, east MP where winter rainfall is mostly due to western disturbances is also decreasing significantly. The decreasing pre- monsoon rainfall over most parts of the central India. The study concluded that the convective activity, which is the main cause for the rainfall activities

during the pre- monsoon season, is decreasing over the central parts of the country. Rainfall is decreasing significantly for the six subdivisions viz. Gujarat Region, West M.P., East M.P., Vidarbha, Chhattisgarh and Jharkhand. Goswami *et al.* (2006) ^[7] visualized the negative impacts of rising global surface temperature and stability of the Indian monsoon rainfall over the past century and investigated the reasons behind this using a daily rainfall data set, and found significant rising trends in the frequency and the magnitude of extreme rain events and a significant decreasing trend in the frequency of moderate events over central India during the monsoon seasons from 1951 to 2000. The seasonal mean rainfall does not show a significant trend because of contribution from heavy events and decreasing moderate events and expected substantial increase in hazards related to heavy rain over central India in the future. Vijay kumar *et al.* (2010) ^[11] observed no significant trend for annual, seasonal, and monthly rainfall for the country as a whole. Annual and monsoon rainfall decreased, while pre-monsoon, post-monsoon and winter rainfall increased at national scale. Rainfall in June, July and September decreased, whereas August rainfall increased, at the national scale. Singh and Mal (2014) ^[25] studied rainfall trends and variability over western Himalaya and found declining trend in annual and monsoon rainfall over high altitudes and increasing over low altitude. Agarwal (1952) ^[1] investigated the trends in rainfall over central India and found a steady change which was not significant during 1908-1940. Kothiyari *et al.* (1997) ^[10], analysed for evaluation of changes in rainfall and temperature regimes in the upper and middle parts of the Ganga basin in northern India. The analysis shows that the total monsoon rainfall and the number of rainy days during the monsoon season are on the decline whereas the annual maximum temperature is on the rise. These changes are observed to have begun around the second half of the 1960's. The results point towards a possible change in the climatic regime. Parthasrathy and Dhar (1976) ^[37] observed increasing trend for annual rainfall over East MP and West MP for 1901-1960 periods.

Indian summer monsoon rainfall and associated large scale circulation are mainly controlled by global parameter like ENSO and NAO, which are indicative of quantification of ocean atmosphere coupling in which sea surface temperature also plays influential role. Mooley and Parthasarathy (1983) ^[13], investigated the relationships of indices of dryness and wetness over India with SOI and SST anomalies are expected to be useful in understanding implications of large-scale anomalies in performance of Indian summer monsoon. Rasmusson and Carpenter (1983) ^[21] and Shukla (1987) ^[36] studied the link between the inter-annual variability (IAV) in the Indian summer monsoon rainfall (ISMR) and the El Nino-Southern Oscillation (ENSO) phenomenon. Rajeevan *et al.* (2008) ^[20] examined variability and long-term trends of extreme rainfall events over central India using 104 years (1901-2004) of high resolution daily gridded rainfall data, variability and long-term trends of extreme rainfall events over central India and found inter- annual and inter-decadal variability of extreme rainfall events over India is influenced by the SST variations over the tropical Indian Ocean. The strong coupling between the equatorial Indian Ocean SST and very high rainfall (VHR) events over the central India suggests that in the present global warming scenario, the frequency of VHR

events and risk of floods may increase over the central India. Yadav (2009) [32] studied inter-annual variability of the Indian summer monsoon rainfall (ISMR) in association with the variability surface temperature, the mean sea level pressure (MSLP), the tropospheric geopotential height and wind patterns over the globe for 1949-2005 period and found the El Nino-Southern Oscillation (ENSO) relationship over ISMR has weakened while the northwest (NW) of North Atlantic sea surface temperature (SST) relationship has increased recent decades. Yadav (2008) [33] established link between 2mST over tropical southwest Pacific and intensification of subtropical westerly jet over Indian Ocean and enhancement of monsoon circulation during negative phase of ENSO. While during non ENSO year positive relationship of ISMR with 2mST anomaly over northwest of North Atlantic Ocean. Srivastava *et al.* (2020) [26] observed ENSO impact on July and August rainfall and found significant changes at multidecadal time scale. The impact of ENSO was significantly strong in August and significantly weak in July during 1948-1980 and post 1980s, ENSO monsoon relationship weak in August than in July.

The present study investigates possible changing influence of southern Oscillation Index (SOI) and North Atlantic oscillation Index (NAO) on rainfall trends over MP.

2. Study area

Madhya Pradesh is also known as the "heart of India", owing to its central location is the second largest state in the country. It is surrounded by five states like Rajasthan, Chhattisgarh, Uttar Pradesh, Maharashtra and Gujarat, completely land locked. It is located between latitude 21.2°N-26.87°N and longitude 74°59' - 82°06'E (figure 1a*). MP has a highly variable subtropical climate also influenced by presence of tropic of cancer which passing through 14 districts. According to Koppen and Geiger climate classification, the state has three main categories i.e. hot summer Mediterranean climate (Csa), tropical savana climate (Aw) and hot semi-arid climate (Bsh). Hot dry summer, followed by monsoon rains (July–September) and a cool and relatively dry winter are the main features of its climate.




Fig 1a: Location map of the study area.

3. Data Used

The data of monthly rainfall (mm) over East MP and West MP meteorological subdivisions from 1871 to 2016 is obtained from Indian Institute of Tropical Meteorological (IITM) Indian regional/sub divisional Monthly Rainfall data set (IITM-IMR). This data series constructed by taking subdivision wise monthly rainfall data of IMD (Primary Data Source). SOI and NAO monthly data obtained from <http://www.ncdc.noaa.gov>.

3. Methodology

Rainfall characteristics such as mean, standard deviation (SD), coefficient of variation (CV) and percentage contribution to annual computed for monthly and season-wise viz., Pre-monsoon (March–May), Monsoon (June–September), Post-monsoon (October–November) and Winter (December–February) for East MP and West MP meteorological subdivision and MP as a whole and compared with all India. The non-parametric Mann-Kendall test along with Sen's slope estimator are used for detection

of trends in rainfall time series 1871-2016 and also to understand recent 50 years trend, considered 1957-2016 time series separately. The seasonal values of Southern Oscillation Index (SOI) and North Atlantic Oscillation (NAO) are prepared from monthly values and the correlation with seasonal and annual rainfall has been calculated.

3.1. Trend Analysis

3.1.1 Non-Parametric -Mann Kendall test for trend analysis

The parametric techniques generally used for assessing central tendency, linear relationship and change point identification. The parametric techniques are based on assumptions of homogeneity and normality. The parametric methods do not perform satisfactorily when there are outliers in the data sets and also if data sets have observations reported as being "less than" or "greater than" some specific values. Parametric methods require substituting some numerical value for such data points will

lead to inexact estimate. Atmospheric variable like precipitation exhibit a marked skewness partly due to the influence of natural phenomena, and hence do not follow a normal distribution, (Mukherjee *et al.*, 2014) ^[15]. The non-parametric techniques are distribution free and can handle presence of outliers in data series and can be used when data are non-normal (Lanzante, 1996) ^[12]. The Mann Kendall test derived by use of a rank correlation test for two groups of observations. If there is positive serial correlation in time series it may lead to significant trend detection without any actual trend (Cox and Stuart, 1955) ^[6].

Mann Kendall test (Mann, 1945) ^[14] a rank-based non-parametric method that has been used to detect trends in hydrological and meteorological time series such as water quality data in lakes and streams (Zetterqvist, 1991; Bouchard and Haemmerli, 1992) ^[34, 3], stream flow data (Chiew and McMahon, 1993; Burn, 1994) ^[5, 4] and meteorological parameter like rainfall and temperature data (Shukla *et al.*, 2017) ^[24]. Detail description or reference list of the Mann -Kendall trend test provided by (Hirsch *et al.*, 1982) ^[9].

The behaviour of annual, seasonal and monthly time series of rainfall is studied by subjecting them to non-parametric Mann-Kendall test and increasing or decreasing slope of trends in the time series is determined by using Sen's method (Sen, 1968) ^[22]. The statistically significant trend detected by use of MK test can be complemented with Sen's slope estimation for magnitude of the trend. The test has carried out with the hypothesis (H₀) that there is no trend in the rainfall time series but it has an alternative hypothesis (H_A) that there is monotonic trend in rainfall time series. For hypothesis testing, the significance levels 0.05(*), 0.01 (* *), and 0.1(+) used for rejecting null hypothesis. The Sen's Slope estimator is a nonparametric, linear slope estimator that works on monotonic data. It is not affected by gross data errors, outliers, or missing data. The Sen's Slope technique is used to determine the magnitude of the trend line. The approach involves computing slopes for all the pairs of ordinal time points using the median of these slopes as an estimate of the overall slope. The calculations are done through the equations used by researchers (Patra *et al.*, 2012) ^[18]. The positive value of test statistic indicates positive trend and negative value indicates negative trend and Sen's slope further quantify the trend.

4. Results and Discussion

4.1 Characteristics of Rainfall

The rainfall characteristics of East MP, West MP and MP as a whole for the period of 1871–2016 and 1967-2016 are calculated. The mean annual rainfall of MP is 1102.0 mm with a standard deviation of 177.8 mm, coefficient of variance 16.1%, with monsoon contribution 90.1% for the period 1871-2016, whereas all India mean annual rainfall 10859 mm with standard deviation 1010 mm and coefficient

of variance 9.3% with monsoon contribution 78.1%. The mean annual rainfall of MP, 1078.3 mm with a standard deviation of 171.1mm, coefficient of variation is 15.9% with monsoon season contribution of 89.7% for the period 1967-2016, whereas all India mean annual rainfall 10700.1 mm with a standard deviation 942.07 mm with monsoon contribution 77.8%. East MP subdivision, the mean annual rainfall is 1249.6 mm having standard deviation of 212.7 mm with coefficient of variance 17% and monsoon contribution 88.9% to annual rainfall for the period 1871-2016 and 1194.6 mm mean annual rainfall with a standard deviation of 214.0 mm, coefficient of variation is 17.9%, monsoon contributes 88.5% to the mean annual rainfall for period 1967-2016. West MP subdivision, the mean annual rainfall is 954.5 mm with a standard deviation of 176.5 mm, coefficient of variance 18.5%, with monsoon contribution 91.2% for the period 1871-2016 and 962.1 mm mean annual rainfall with a standard deviation of 162.9 mm, coefficient of variation is 16.9%, monsoon season contributes 91.3% to the mean annual rainfall for period 1967-2016.

July and August month rainfall amount is very crucial for monsoon as about 64 to 66% of seasonal rainfall occur in these months for East MP and West MP subdivisions. However spatial and temporal daily rainfall distribution is very much variable over the state leading to flood and drought situations in different parts of the state, where as for all India, the mean rainfall of June, July, August and September is 163.1, 272.5, 242.2 and 170.3 mm respectively for 1871-2016. The rainfall of July month contributes 25% while August rainfall contributes 22.3% to the annual rainfall. The June and September rainfall contribute 15.1% and 15.7% to the annual rainfall respectively. The mean seasonal rainfall are 23.2 mm, 94.4 mm, 848.1 mm and 120.1 mm respectively.

4.2 Annual Trend

The trends are detected using non parametric Mann Kendall test and presented in Table 1 and Table 2. The statistical techniques are the way to generalize the results form a large data set and significance testing assures that the results are not by chance. The mean annual rainfall over MP showed a declining trend (Fig 2a) which is not significant while trends for West MP increasing but not significant (Fig.2b). There is decrease of 55 mm mean rainfall compared to long-term mean annual rainfall for MP as a whole. The long-term trends for East MP subdivision are decreasing significantly at 0.05 level. The declining trends are more prominent from 1941 onwards and during recent decade a decrease of 143 mm of mean annual rainfall with reference to long-term mean of annual rainfall for the subdivision (Fig 2c). The decreasing trends over East MP subdivision have been indicated by many other researchers also while all India annual rainfall showing positive trend but not significant.





Fig 2a: Annual rainfall trend for MP, 2b: West MP, 2c: East MP

4.3 Seasonal Trends

4.3.1 Pre-monsoon (March – May)

The pre-monsoon rainfall has an increasing trend for MP state, West MP and East MP subdivision. The ten years moving average showing increasing signature from 1972 for MP (Fig.2a) from 1960 for West MP subdivision (Fig.2b) and from 1972 for East MP subdivision (Fig.2c). There is

increase of 2.47 mm in past ten year mean pre-monsoon rainfall for MP as compared to long-term mean rainfall for pre-monsoon. These trends are not significant. The all India pre-monsoon rainfall showing negative trend for whole 1871-2016 period while positive trend for recent fifty years period (1967-2016) but not significant (table 1 and table 2).






Fig 2: Pre-monsoon rainfall trend for MP (2a), West MP (2b), East MP (2c)

4.3.2 Monsoon (June -September)

The monsoon is very important phenomena for economy and agriculture of the state. The land lock central state fully depends on monsoon for its water and associated requirements. The long-term monsoon trends are decreasing at 0.1 level of significance for the MP state (Fig.3a). The observed trends are more or less constant over west MP subdivision (Fig.3b) while the trends are significantly decreasing for East MP at 0.01 level (Fig.3c). There is

decrease of 99.147 mm and 196.15 mm in past ten year mean monsoon rainfall for EMP and MP as a whole compared to long-term mean rainfall for monsoon. The uneven spatial distribution of rainfall in monsoon and year to year variability are precursors of climate change over the state. The all India monsoon rainfall showing non-significant positive trends for 1871-2016 and 1967-2016 period (Table 1 and Table 2).




Fig 3: Monsoon rainfall trend for MP (3a), West MP (3b), East MP (3c)

4.3.3 Post monsoon (October November, December)

Post- monsoon contributes 5% to annual rainfall and trends

are decreasing over East MP subdivision while long trends are over MP and West MP subdivision are non-significant,

during past fifty (1967-2016) years trend are significantly decreasing for MP state at 0.05 level (Fig.4a), 0.1 level over West MP (Fig.4b) and no significant trend in East MP

(Fig.4c). The all India post-monsoon trends are negative for the whole 1871-2016 period.





Fig 4: Post-monsoon rainfall trend for MP (4a), West MP (4b), East MP (4c)

4.3.4 Winter (January-February)

The long-term trends for winter season for West MP, East MP and MP state are nearly constant (Fig. 5a, 5b, 5c). The winter season contributes 2.5 to annual rainfall for the state and it is mainly due to influencing western disturbances.

The frequency of western disturbances and penetration of trough to lower latitude contributes good rainfall during the season. The all India winter rainfall trends are negative for 1871-2016 and 1967-2016 period but not significant.






Fig 5: Winter rainfall trend for MP (5a), West MP (5b), East MP (5c)

4.3.5 Monthly rainfall trends

Monthly rainfall trends for all months have been observed by Mann-Kendall test and time series also fragmented into two parts to identify recent monthly trends. The results are shown in Table 1 and Table 2. The Long-term (1871-2016) June month rainfall is significantly decreasing over East MP, West MP and MP as a whole state, July rainfall tends decreasing over East MP and significant at 0.05 level and August rainfall trends are significantly increasing at level 0.05 over West MP. February rainfall trends are decreasing for both subdivisions and the state as a whole. November and December month almost constant over the study period. For all India, June and July rainfall trends are decreasing significantly at 0.1 level. If we consider recent past fifty years (1967- 2016), July rainfall trends are increasing over West MP and MP state significantly while August month rainfall is showing decreasing trends at 0.05 level of significance over East MP and 0.1 level of significance over West MP. The rainfall over the state in August is significantly decreasing at 0.05 level of significance. September and December rainfall trends are decreasing over the state. The trends are not significant in rest months for the state while monthly trends are not significant for all India in recent fifty years.

4.3.6 Decadal variations in annual and seasonal rainfall

The decadal mean and its departure from normal for annual, seasonal rainfall and frequency of excess and deficient years are presented in (Table 3, Table 4, and Table 5). The rainfall greater than one standard deviation from mean is termed as excess rainfall and less than one standard deviation from mean is termed as deficit rainfall. West MP subdivision sharing boundaries with Rajasthan, Gujarat and Maharashtra decadal mean rainfall during (1871-1880) in monsoon season, four excess and one deficient and during six decades (1951-2010) five excess and nine deficient years observed. Annual rainfall over West MP during (1871-1900), three excess and three deficient and during (1901-1930), three excess and ten deficient years were observed. During 1871-2016, excess rainfall years are greater than deficient rainfall years for winter, pre-monsoon and post monsoon while deficient years are greater than excess rainfall years for

monsoon season and also on annual basis.

East MP subdivision touching boundaries of Uttar Pradesh, Chhattisgarh and Vidarbha region, the decadal mean monsoon rainfall during three decades (1871-1900), seven excess and three deficient and during recent six decades (1951-2010) only five excess and nine deficient years were observed. Trend of excess monsoon rainfall years are decreasing. No excess year was observed for post- monsoon season during (1951-2010). No deficient and no excess years for winter season during (1931-1960). During (1871-2016), twenty-four excess and twenty-one deficient years are there on annual basis. Over the MP state as a whole during initial five decades (1911-1960), twelve excess and six deficient years were noticed while recent five decades (1961-2010) five excess and eleven deficient years for monsoon and five excess and seven deficient years for post-monsoon season are there. This again confirms the decreasing post-monsoon rainfall over the state in recent fifty years. The total numbers of excess rainfall years are greater than deficient years for annual and seasonal basis. Monsoon season is mainly rain giving season for MP and excess and deficient rainfall frequencies seem to have some signals of fifty years transition period for decadal deficient years to decadal excess year.

5. Tele-connections

The Southern Oscillation Index (SOI) is a measure of strength of the walker circulation and also indicative of El Niño and La Niña events and impact on Indian monsoon. The SOI measures the difference in surface air pressure between Tahiti and Darwin. The correlation between Indian monsoon rainfall and Southern Oscillation Index (SOI) was first explained by Walker (1923-1924) [28, 29]. The Southern Oscillation Index (SOI) is the atmospheric component of ENSO. The studies in recent years showed strong inter-decadal variability in the ENSO cycle and due to this, state of Pacific Ocean and its influence on monsoon is also variable on decadal scale (Webster *et al.*1998) [30]. Various research studies have done to explore relationship between rainfall and oceanic index SOI.

The rainfall trends and the possible link with of SOI and NAO are investigated (Fig.6 and Fig.7) for annual and

seasonal rainfall over the state. SOI has positive correlation with annual and seasonal rainfall over East MP and west

MP subdivisions and MP except winter season for 1876-2016 period.




Fig 6: SOI values for 1876-2016 and 1967-2016 for MP, WMP and EMP

The correlation of annual and monsoon rainfall with SOI significant at 0.05 level for West MP and MP state while NAO has significant positive correlation with monsoon rainfall over East MP and MP as a whole at 0.05 level on long-term basis. During recent fifty years 1967-2016, the SOI relationship has weakened for annual and monsoon rainfall over West MP, East MP and over the state. Winter

season rainfall is negatively correlated with SOI and NAO during whole 147 years. In contrast to this post monsoon season rainfall correlation became negative with SOI during recent fifty years while remained negative with NAO for the whole period. The transition from positive to negative correlation of SOI explains decreasing post-monsoon rainfall in recent fifty years over the state.




Fig 7: NAO values for 1876-2016 and 1967-2016 for MP, WMP and EMP

Table 1: Rainfall trends over All India, M.P., East M.P., and West M.P. for 1871-2016

Months	All India		MP		East MP		West MP	
	Sen's Slope	Test Z	Sen's Slope	Test Z	Sen's Slope	Test Z	Sen's Slope	Test Z
Jan	-0.09	-0.69	0.00	0.40	0.01	0.73	0.00	-0.27
Feb	-0.05	-0.42	0.00	-0.17	0.00	-0.13	0.00	-0.30
Mar	0.05	0.32	0.01	1.45	0.01	1.04	0.00	1.08
Apr	0.26	1.57	0.01	1.38	0.01	1.19	0.00	1.29
May	0.06	0.18	0.00	0.24	0.01	0.53	0.00	0.04
Jun	-1.37	-1.66	-0.37	-2.32*	-0.42	-2.39*	-0.28	2.06*
Jul	-1.07	-1.69	-0.22	-1.14	-0.60	-2.57*	0.15	0.83
Aug	0.71	0.89	0.22	1.19	-0.01	-0.05	0.42	2.12*
Sep	-0.67	-0.92	-0.23	-1.30	-0.23	-1.23	-0.23	-1.21
Oct	0.32	0.59	0.03	0.85	0.02	0.51	0.02	1.14
Nov	0.04	0.13	0.00	0.25	0.00	-0.16	0.00	0.98
Dec	-0.03	-0.26	0.00	-1.16	0.00	-0.15	0.00	-0.59
Winter	-1.05	-0.58	0.01	0.32	0.03	0.57	0.01	0.20
Pre-monsoon	-0.13	-0.58	0.03	0.95	0.04	0.80	0.02	0.80
Monsoon	0.28	0.74	-0.66	-1.84+	-1.24	-2.98**	-0.09	-0.21
Post-monsoon	-2.06	-1.28	0.02	0.31	0.00	-0.04	0.04	0.64
Annual	0.41	0.57	-0.52	-1.42	-1.13	-2.53*	0.12	0.26

Legends: - + : 0.1 , * : 0.05, ** : 0.01

Table 2: Rainfall trends over All India, M.P., East M.P., and West M.P. for 1967-2016

Months	All India		MP		East MP		West MP	
	Sen's Slope	Test Z	Sen's Slope	Test Z	Sen's Slope	Test Z	Sen's Slope	Test Z
Jan	-0.18	-0.39	-0.02	-0.26	-0.06	-0.74	0.01	0.49
Feb	-0.07	-0.17	-0.03	-0.34	-0.07	-0.61	0.00	-0.34
Mar	0.00	0.00	0.00	0.03	0.01	0.22	0.01	0.23
Apr	0.70	0.75	0.04	1.05	0.07	1.22	0.00	0.25
May	1.53	1.10	0.03	0.23	0.05	0.61	0.00	0.05
Jun	2.20	0.62	0.04	0.04	0.30	0.33	-0.13	-0.23
Jul	-2.07	-0.57	1.23	1.66+	0.52	0.49	1.81	1.90+
Aug	-4.46	-1.27	-2.18	-2.20*	-2.69	-2.26*	-1.67	-1.73+
Sep	2.24	0.66	-0.70	-1.00	-0.85	-1.01	-0.59	-0.61
Oct	0.22	0.13	0.11	0.57	0.18	0.69	0.02	0.16
Nov	-0.25	-0.28	0.00	0.29	0.00	0.22	0.00	-0.12
Dec	-0.61	-1.02	-0.01	-1.15	0.00	-0.97	0.00	-0.94
Winter	-2.86	-0.33	0.00	0.05	0.02	-0.43	0.03	0.76
Pre-monsoon	0.31	0.31	0.01	0.27	0.21	0.50	-0.03	-0.46
Monsoon	2.20	1.04	-0.01	-1.19	0.01	-1.31	0.03	-0.47
Post-monsoon	-4.80	-0.79	-1.19	-2.01*	0.01	-1.40	-0.58	-1.80+
Annual	-1.91	-0.66	-0.53	-1.24	-0.51	-1.20	0.04	-0.42

Legends: - + : 0.1 , * : 0.05, ** : 0.01

Table 3: Decadal mean (departure from normal), frequency of excess and deficit years for M.P.

Decade	Annual			Pre-Monsoon			Monsoon			Post-Monsoon			Winter		
	Decadal mean	Excess	Deficit	Decadal mean	Excess	Deficit	Decadal mean	Excess	Deficit	Decadal mean	Excess	Deficit	Decadal mean	Excess	Deficit
1871-1870	3	3	1	-4	1	1	5.6	3	1	-29.9	1	3	-17.7	1	2
1881-1890	7.3	2	0	-16.3	0	2	8	3	0	24.3	3	0	-31.6	0	4
1891-1900	-1	2	2	-26.3	1	3	0.6	1	1	-18.5	1	2	-3.1	1	1
1901-1910	-9.6	0	3	-1.5	1	0	-8.8	0	3	-32.2	1	4	-0.5	2	1
1911-1920	-1.5	3	3	41.4	3	0	-4.1	2	3	14.6	2	1	20.4	1	2
1921-1930	-1	1	0	-11.6	0	1	-1.5	1	0	19.7	2	1	-14.6	0	2
1931-1940	7.9	2	0	-16.6	1	0	7.5	1	0	21.9	2	1	16.6	0	0
1941-1950	9.7	4	1	-8.3	1	0	8.9	5	1	1.8	2	1	-8.3	4	0
1951-1960	0.5	2	2	1.1	0	0	0.1	0	1	11	2	0	-1.1	1	0
1961-1970	-3.3	1	2	-6.5	1	0	-2.0	1	2	-16.2	1	1	-6.5	1	2
1971-1980	1.6	3	2	-21.3	1	2	0.8	2	2	27.7	0	2	-21.3	2	1
1981-1990	-3.7	1	1	2.7	1	0	-5.3	1	2	0.9	2	0	2.7	4	1
1991-2000	-1.7	2	2	12.1	2	0	-2.1	1	2	11.7	1	1	12.1	1	2
2001-2010	-12.3	0	5	4.5	2	1	-12.3	0	3	-13.7	1	1	4.5	1	1
2010-2016	6.9	1	0	30.6	1	0	7.6	2	1	-36.1	0	2	30.6	2	1

Table 4: Decadal Mean (departure from normal), frequency of excess and deficit years for West M.P.

Decade	Annual			Pre-monsoon			Monsoon			Post-monsoon			Winter		
	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit
1871-1880	4.6	1	1	15.6	0	0	6.8	4	1	-27.9	0	3	-2.1	2	2
1881-1890	4.7	2	0	-7.8	0	0	4.9	2	1	16.2	3	0	-26.3	1	1
1891-1900	-4.6	0	2	-24	1	2	-2.8	0	2	-30.6	2	1	-4.2	1	1
1901-1910	-14.6	0	4	-13.4	1	0	-13	0	4	-46.7	2	2	-8.7	2	0
1911-1920	-7.3	3	4	52	2	0	-10.3	2	4	13.3	2	1	32.2	0	3
1921-1930	-83	0	2	-35.7	0	1	-9.1	0	2	15.3	3	1	3.8	1	1
1931-1940	9.3	3	0	-18.3	0	0	9.3	2	0	24.3	2	0	-48	0	0
1941-1950	13.6	4	1	-12.6	1	0	13.8	4	0	3.8	3	1	63.3	3	0
1951-1960	13	2	1	5.9	1	0	0.2	1	1	18.41	0	0	6.1	0	0
1961-1970	-0.5	1	2	-13.8	1	0	0.8	1	2	-11.4	0	1	-25.8	0	2
1971-1980	5	1	1	-20	0	0	5	1	1	23.3	0	0	-18.8	1	0
1981-1990	-2.1	0	2	12.9	1	1	-4.3	0	1	24.9	0	0	26.2	3	0
1991-2000	0.7	1	3	29.1	1	0	-0.7	1	1	23.5	0	1	-19.6	1	0
2001-2010	-10.5	1	0	12.8	1	0	-10.5	1	3	-13.5	0	0	-23.3.	0	1
2010-2016	14.4	1	0	21.2	1	0	16.6	2	0	-54.9	1	2	88.1	3	0

Table 5: Decadal mean (departure from normal), frequency of excess and deficit years for East M.P.

Decade	Annual			Pre-monsoon			Monsoon			Post-monsoon			Winter		
	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit	Decadal Mean	Excess	Deficit
1871–1880	1.7	2	0	8.7	1	1	4.6	2	1	-31.5	0	0	-24.2	1	2
1881–1890	9.2	2	0	-2.04	0	1	10.5	2	1	29.9	3	0	-33.8	1	4
1891–1900	1.8	2	1	28.1	1	3	3.3	3	1	-9.8	2	3	5.8	1	1
1901–1910	3	0	2	4.5	0	1	-5.5	1	2	-21.6	2	3	2.6	2	0
1911–1920	4.5	3	3	36.2	2	0	0.8	2	3	15.3	2	1	15.5	1	2
1921–1930	6.9	2	0	0.5	1	2	4.5	2	1	22.6	3	1	-19.1	0	2
1931–1940	6.8	2	0	15.8	1	0	6.1	1	0	19.9	2	1	1.5	0	0
1941–1950	-0.1	1	1	-6	1	0	5.1	2	1	0.1	3	1	74.3	0	0
1951–1960	-5.5	2	1	-1.2	2	0	0	1	0	5.4	0	1	-8.6	3	0
1961–1970	-1.1	1	2	-21.9	1	0	-4.2	1	2	-19.9	0	1	-21.3	2	1
1971–1980	-50	2	3	-2.4	0	2	-6.1	1	3	30.7	0	0	-5.3	4	1
1981–1990	-3.5	1	1	3.6	0	0	-3.2	1	2	-16.7	0	0	44.3	1	2
1991–2000	-13.8	2	2	3.6	0	0	-13.6	1	1	-0.8	0	2	-27.5	1	1
2001–2010	1.2	0	5	35.4	0	1	0.6	0	2	-14	0	1	-29.6	1	1
2010–2016	1.6	2	0	21.3	1	0	0.6	2	2	-22.4	1	1	24.4	1	1

Conclusions

The long-term monsoon rainfall trends are significantly decreasing over MP and mainly this decrease is more prominent over East MP. June month rainfall is significantly decreased over MP. A significant decrease in July rainfall over East MP and August rainfall over West MP. June month rainfall mainly depends up on timely onset of monsoon and meso-scale convective activities in the state. The delay in monsoon over the state lead to less rainfall in June and also affect sowing pattern of kharif crops in mainly in rainfed area. During recent fifty years the trends for the post monsoon rainfall are decreasing at 0.05 level of significance over the state MP and monthly rainfall trends for August are decreasing over East MP at 0.05 level of significance and significant decrease over West MP also at 0.1 level. July month rainfall trends are increasing over west MP and MP at 0.1 level of significance. The post monsoon season rainfall is crucial for maintenance of soil moisture for sowing of Rabi crops in the state. This decrease in rainfall will lead to more dependence on ground and surface water resources for irrigation in the state which leads to increase stress on water resources.

The ground water is the source of drinking water and decrease in rainfall leads to decrease in ground recharge and over exploitation of ground water during summer season (irrigation, industry and drinking water). The tele-connections have given indication of weakening of SOI correlation with monsoon rainfall and the positive correlation turned into negative correlation during past fifty years for post monsoon season. Studies have shown positive relationship between SOI and south west monsoon rainfall during monsoon season starts building during March to May and turns strong in concurrent mode.

References

1. Agarwal KS. Fluctuations of annual rainfall in central India 1872-1947, *Indian J Meteor Geophys.* 1952;3(3):229-230.
2. Barnard GA. Significance tests for 2x2 tables, *Biometrika.* 1947;34(1/2):123. doi:10.2307/2332517.
3. Bouchard A, Haemmerli J. Trend detection in water quality time series of LRTAP-quebec network lakes, *Water, Air, & Soil Pollution.* 1992;62(1-2):89-110. Doi:10.1007/bf00478455.
4. Burn DH. Hydrologic effects of climatic change in

west-central Canada, *Journal of Hydrology.* 1994;160(1-4):53-70. Doi:10.1016/0022-1694(94)90033-7.

5. Chiew FHS, McMahon TA. Detection of trend or change in annual flow of Australian rivers, *International Journal of Climatology.* 1993;13(6):643-653. Doi:10.1002/joc.3370130605.
6. Cox DR, Stuart A. Some quick sign tests for trend In location and dispersion, *Biometrika.* 1955;42(1/2):80. Doi:10.2307/2333424
7. Goswami BN, Venugopal V, Sengupta D, Madhusoodanan MS, Xavier PK. Increasing trend of extreme rain events over India in a warming environment, *Science.* 2006;314(5804):1442-1445. Doi:10.1126/science.1132027.
8. Guhathakurta P, Rajeevan M. Trends in the rainfall pattern over India, *International Journal of Climatology.* 2008;28(11):1453-1469. Doi:10.1002/joc.1640.
9. Hirsch RM, Slack JR, Smith RA. Techniques of trend analysis for monthly water quality data, *Water Resources Research.* 1982;18(1):107-121. Doi:10.1029/wr018i001p00107
10. Kothiyari UC, Singh VP, Aravamuthan V. Journal search results-cite this for me, *Water Resources Management.* 1997;11(1):17-34. Doi:10.1023/a:1017936123283
11. Kumar V, Jain SK, Singh Y. Analysis of long-term rainfall trends In India, *Hydrological Sciences Journal.* 2010;55(4):484-496. Doi:10.1080/02626667.2010.481373
12. Lanzante JR. Resistant, robust and non-parametric techniques for the analysis of climate data: Theory and examples including application to historical radiosonde station data, *International Journal of Climatology.* 1996;16(11):1197-1226. Doi:10.1002/(sici)1097-0088(199611)16:11<1197::aid-joc89>3.0.co;2-1.
13. Mooley DA, Parthasarathy B. Variability of the Indian Summer Monsoon and Tropical Circulation Features, *Monthly Weather Review.* 1983;111(5):967-978. Doi:10.1175/1520-0493(1983)111<0967:votism>2.0.co;2.
14. Mann HB. Nonparametric tests against trend, *Econometrica.* 1945;13(3):245. Doi:10.2307/1907187.
15. Mukherjee S, Joshi R, Prasad RC, Vishvakarma SCR, Kumar K. Erratum to: summer monsoon rainfall trends

- in the Indian Himalayan region, *Theoretical And Applied Climatology*. 2014;121(3-4):803-805. Doi:10.1007/s00704-014-1319-4.
16. Naidu CV, Srinivasa Rao BR, Bhaskar Rao DV. Climatic trends and periodicities of annual rainfall over India, *Meteorological Applications*. 1999;6(4):395-404. Doi:10.1017/s1350482799001358
 17. Parthasarathy B, Dhar ON. Study of trends and periodicities of rainfall over M.P, *Proc. Ind. National Sc. Academy*. 1976;42-A(1):73-80.
 18. Patra JP, Mishra A, Singh R, Raghuvanshi NS. Detecting rainfall trends in twentieth century (1871–2006) over Orissa State, India, *Climatic Change*. 2011;111(3-4):801-817. Doi:10.1007/s10584-011-0215-5.
 19. Radhakrishnan K, Sivaraman I, Jena SK, Sarkar S, Adhikari S. A Climate Trend Analysis of Temperature and Rainfall in India, *Climate Change And Environmental Sustainability*. 2017;5(2):146. Doi:10.5958/2320-642x.2017.00014.x.
 20. Rajeevan M, Bhatte J, Jaswal AK. Analysis of variability and trends of Extreme Rainfall Events Over India Using 104 Years Of Gridded Daily Rainfall Data, *Geophysical Research Letters*. 2008, 35(23). Doi:10.1029/2008gl036105
 21. Rasmusson EM, Thomas HC. The relationship between eastern equatorial Pacific sea surface temperatures and rainfall over India and Sri Lanka, *Monthly Weather Review*. 1983;111(3):517-528. Doi:10.1175/1520-0493(1983)111<0517:trbeep>2.0.co;2.
 22. Sen PK. Estimates of the regression coefficient based on Kendall's tau, *Journal of The American Statistical Association*. 1968;63(324):1379-1389. Doi:10.1080/01621459.1968.10480934
 23. Sharma HO, Rathi D, Chauhan RS, Meena SC. Status of Agriculture in Madhya Pradesh, A report submitted to MoA, GoI, New Delhi; c2015.
 24. Shukla R, Khare D, Tiwari P, Mishra P, Gupta S. Analysis of long-term temperature trend for Madhya Pradesh, India (1901-2005), *Current World Environment*. 2017;12(1):68-79. doi:10.12944/cwe.12.1.09.
 25. Singh RB, Mal S. Trends and Variability of Monsoon and Other Rainfall Seasons in Western Himalaya, India, *Atmospheric Science Letters*. 2014;15(3):218-226. doi:10.1002/asl2.494.
 26. Srivastava G, Chakraborty A, Nanjundiah RS. Multidecadal variations In ENSO-Indian Summer Monsoon relationship at sub-seasonal timescales, *Theoretical and Applied Climatology*. 2020;140(3-4):1299-1314. Doi:10.1007/s00704-020-03122-6.
 27. Sekhar PH, Dr. Kesavulu Poola, Sekhar KR, Dr. Naidu MB. Modelling and prediction of coastal Andhra rainfall using ARIMA and ANN models. *Int. J Stat Appl Math* 2020;5(6):104-110.
 28. Walker GT. Correlation in seasonal variations of weather, VIII: A preliminary study of world weather, *Mem. India Meteor. Dep*; 1923;24:75-131.
 29. Walker GT. Correlation in seasonal variations of weather, IX: A further study of world weather, *Mem. India Meteor. Dep*. 1924;24:275-332
 30. Webster PJ, Magana VO, Palmer TN, Shukla J, Tomas RA, Yanai M, *et al.* Monsoons: processes, predictability, and the prospects for prediction, *Journal Of Geophysical Research: Oceans*. 1998;103(C7):14451-14510. doi:10.1029/97jc02719
 31. Xu Z, Liu Z, Fu G, Chen J. Trends of major hydroclimatic variables In the Tarim river basin during the past 50 years, *Journal Of Arid Environments*. 2010;74(2):256-267. Doi:10.1016/j.jaridenv.2009.08.014.
 32. Yadav RK. Changes In The large-scale features associated with the Indian Summer Monsoon in the recent decades, *International Journal Of Climatology*. 2009;29(1):117-133. doi:10.1002/joc.1698.
 33. Yadav RK. Role of equatorial Central Pacific and northwest of North Atlantic 2-metre surface temperatures in modulating Indian summer monsoon variability, *Climate Dynamics*. 2008;32(4):549-563. Doi:10.1007/s00382-008-0410-
 34. Zetterqvist L. Statistical Estimation and Interpretation Of Trends In Water Quality Time Series, *Water Resources Research*. 1991;27(7):1637-1648. doi:10.1029/91wr00478.
 35. Kumar S, Rosenberg JM, Bouzida D, Swendsen RH, Kollman PA. The weighted histogram analysis method for free-energy calculations on biomolecules. I. The method. *Journal of computational chemistry*. 1992 Oct;13(8):1011-21.
 36. Shukla J. Interannual variability of monsoons. *Monsoons*. 1987 Feb 23;14:399-464.
 37. Parthasarathy B, Dhar ON. A study of trends and periodicities in the seasonal and annual rainfall of India. *MAUSAM*. 1976 Jan 1;27(1):23-8.
 38. Walker GT. On the meteorological evidence for supposed changes of climate in India, *memoris, Me. Dept*. 1910;21(part 1).