



P-ISSN: 2706-7483  
 E-ISSN: 2706-7491  
 IJGGE 2024; 6(1): 351-355  
<https://www.geojournal.net>  
 Received: 11-11-2023  
 Accepted: 17-12-2023

**Karrar AH Arab**  
 Department of Ecology,  
 College of Science, University  
 of Kufa, Najaf, Iraq

## Evaluation of dustfall rates on Najaf and its temporal and spatial variations

**Karrar AH Arab**

DOI: <https://doi.org/10.22271/27067483.2024.v6.i1e.239>

### Abstract

The increasing problem of air pollution, along with the problem of lack of rainfall, the increase in the area of the dry and desertified area, and the fragmentation of soil, contribute greatly to increasing concern about the environmental damage resulting from these problems. This study aims to examine dustfall in four sites from Najaf province for eleven months, starting in February and ending in December 2021. The direct traditional method using aluminium-collector plate was used to collect falling dust. The results of quantitative analysis of falling dust rates showed that falling dust rates for the study sites during the study period ranged between (110.14 - 1952.95) mg/m<sup>2</sup>/day. The dustfall rate values exceeded the permissible limits set by Australian and Malaysian standards in all the study areas throughout the study period, except two data points. The results also showed a spatial and temporal change in the values of falling dust rates. Falling dust rates indicate the presence of processes or activities that emit high levels of dust. Therefore, these operations or activities should be identified and stopped, and legal action should be taken to prevent these activities from occurring.

**Keywords:** Najaf, Dustfall rate, aluminium-collector plate

### Introduction

The dust is considered as a nuisance when it accumulates and falls, it quickly causes damage to the roof of residential properties or commercial buildings. However, the level at which it becomes a nuisance is very subjective so a nuisance rating may be left up to personal opinions. Experimental data from dustfall quantity measurements are frequently used to assess the level of impact the dust has on the environment <sup>[1]</sup>. Falling dust in meters allow measuring the average dustfall rate, which can be compared to threshold values. The rate of dustfall is expressed as average milligrams per square meter per day (mg/m<sup>2</sup>/day). The average rate of dust falling varies depending on the characteristics of the area. For example, the average rate of dustfall in the city and residential areas differs from that in commercial centres <sup>[2]</sup>. Threshold limit values of dust fall are maintained in a number of recommended air quality guidelines. The table below shows the values of some of them.

The Plant growth is hampered by dust, and dust clogs pores and reduces photosynthesis activity. While transporting dust, many young plants are lost resulting in decreased productivity. Plant infections also occur due to the palm dust mite pest *Oligonychus afrasiaticus* Meg and so is Acari Tetranychida, which is a severe pest in North Africa, the Near East, and other date palm regions of the world. The dust mite attacks palm fruits in their early growth stages. Adult and nymph pests suck the sap of immature green date fruit, causing severe scarring, deformation, browning, and peeling of the fruit <sup>[13]</sup>.

An accurate understanding of the potential risks of dust depends on knowing the type, size, shape and density of dust particles along with wind speed and direction, rainfall, local topography and hydrogeology, which are parameters that can disperse dust emissions. However, in general, large dust particles (larger than 30 µm), which make up the largest proportion of dust emitted by metalwork, will largely settle within 100 meters of the source. Medium-sized particles of 10 to 30 µm are more likely to travel from 200 to 500 m <sup>[14]</sup>.

The importance of the research: In recent decades at the global level, there is much interest in studying the sources of dust, the model of its transportation, its effects on the environment, and its interaction with the problem of climate change. In addition, basic data on the characteristics of spatial and temporal changes are still lacking, not available in all cities.

**Corresponding Author:**  
**Karrar AH Arab**  
 Department of Ecology,  
 College of Science, University  
 of Kufa, Najaf, Iraq

**Table 1:** The threshold limit values of dustfall as (average dustfall rate mg/m<sup>2</sup>/day)

Country	Average of dustfall rate
Department of Environment Malaysia, Malaysia (3)	133 mg/m <sup>2</sup> /day
US Environmental Protection Agency, American (5)	350 mg/m <sup>2</sup> /day
British Environment Agency, Britain (6)	200 mg/m <sup>2</sup> /day
Australian Department of Environment, Australia (7)	4 g/m <sup>2</sup> / month 133 mg/m <sup>2</sup> /day

**Aim of the study**

The aim of this study is to assess the monthly variation in the dust fall rate and an assessment of the potential dangers of the dust fall rates, in addition to a test of the variation in the rate of dustfall between the studied stations for the period from 2/1/2021 to 1/1/2022.

**Methodology**

Samples of dustfall were collected at the study sites, using metal bottles made of aluminium with diameters equal to one-third of their heights. This method of collection is followed by both the Iraqi Environmental Protection and Improvement Department and the Meteorology and Seismic Monitoring Authority in estimating the amounts of falling dust and other studies. Others tools can also be used to collect falling dust, if available, for example Frisbee Bottle [15].

Dust collection containers were distributed to four locations (residential, industrial, commercial) as follows: The first location was named (Aden District/Industrial), and it is an industrial area located south of the holy city of Najaf. The second sample collection station was named (Kufa / commercial) station, which is located in the Kufa District, east of the city of Najaf. The third sample collection station was named (Manathira / Residential), located in the Manathira District, which is about twenty kilometers south of the holy city of Najaf. The fourth station, which was named (Najaf/residential), is located in the center of the city of Najaf, which is a residential area. Note that the bottles were installed in the elected school site at a height of 1.5 m above the floor level of each site, and the dust collection bottles were changed at the end of each month. The collected dust is placed in a beaker containing distilled water, which then evaporates the water to ensure that the results are only representative of materials of a mineral nature. The weight of falling dust was calculated by subtracting the weight of the empty container (W1) from the weight of the container after collecting dust (W2), which is its weight after it was placed on site. Thirty days after thawing and fumigation.

**Calculating the rate of dustfall**

The rate of dustfall is expressed as the average weight of dust falling on an area of one square meter over thirty days.

The unit of measurement is gram square/ meter/thirty days. The rate of dust falling at the end of each month was calculated according to the following steps:

1. Calculate the weight of dustfall collected in each sample collection container during a month at each study site
2. Weight of dustfall = W2- W1
3. The rate of dustfall is gm \m2 per month = (weight of dust in the container / area of the base of the container in square metres)
4. Calculating the area of the collection container (cm<sup>2</sup>) = radius squared \* π.

**Statistical Analysis**

The data represent in the form of a box, which shows the distribution and spread of data, and through it is possible to determine whether there is extreme (anomalous) data or that is inconsistent with the main data set. As for the method of drawing the box, it is by specifying the first quartile of the data, which is the value that captures less than 25% of the data, and the third quartile, which is the value that captures less than 75% of the data [16].

ANOVA test is a way to find out whether the data collected are similar or different. In other words, it helps to know whether the null hypothesis should be rejected or an alternative hypothesis should be accepted. Basically, you test different groups to see if there is a difference between them. Use one-way analysis of variance One Way ANOVA determines the number of independent variables in the analysis of variance test, the research is one independent variable.

**Results**

The falling dust rate was calculated in mg/m<sup>2</sup>/day. In order to compare the collected data with the values of available standards that specify permissible dust levels in air pollution laws, see table 2. The rate of dust falling to the study sites during the study period ranged between 110.14 - 1952.95 mg/m<sup>2</sup>/day. The average values of the falling dust rate at the four stations at each month are presented in Table 3. The average dust fall rate was calculated for eleven months in each study site and is presented in Table 4. The lowest average values of the falling dust rate were in station 1 (Aden district), which was 482.97 (mg / m<sup>2</sup> / day), while the highest average values of the falling dust rates were in station 4 (Najaf / Residential), which was 612.56 (mg/m<sup>2</sup>/ day) day).

The normal distribution test was conducted for the data of the four stations collected during the study period using the D'Agostino & Pearson normality test and also using the Shapiro- Wilk normality test. The results indicated that the distribution of data from the second and fourth stations was not normal, while the distribution of data from the first and third stations was normal.

**Table 2:** The rate of dustfall in mg/m2/day at four stations in the holy city of Najaf and its suburbs during eleven months in the year 2021

Months	Najaf / Residential	Manathira / Residential	Kufa / Commercial	Aden district/ Industrial
February	256.68	343.82	322.88	347.78
March	288.56	314.98	414.17	553.54
April	272.72	776.28	389.65	172.00
May	1386.40	877.56	424.16	631.62
June	910.94	344.57	387.20	213.68


July	370.98	521.29	306.85	709.51
August	534.12	385.69	1240.61	296.48
September	392.85	425.10	120.14	484.32
October	292.52	1055.41	391.34	246.69
November	305.53	110.14	513.18	198.03
December	386.44	474.71	643.13	649.16

**Table 3:** Average dust fall rate for the study sites in each month

Months	Average rate of dustfall (mg/m <sup>2</sup> /day)
February	317.79
March	392.81
April	402.66
May	829.93
June	464.10
July	477.16
August	614.22
September	355.61
October	496.49
November	281.72
December	538.36

**Table 4:** Average rate of dustfall for eleven months (mg/m<sup>2</sup>/day) at all study sites


Najaf (residential)	Manathira (Residential)	Kufa (commercial)	Aden district (Industrial area)
612.56	557.11	581.03	482.97



**Fig 2:** Values of the rate of dustfall in four sites in Najaf during eleven months from February 2021 to January 2022

It should be noted here that the results of one-way analysis of variance tests using the Kruskal -Wallis test No presence revealed difference statistically significant between the values of the dust falling rate at the stations. It is noted that the highest values of falling dust rates were recorded in the month of May, where the average was 829.9 (mg/m<sup>2</sup>/day). The lowest values of falling dust rates were in the month of November, when the average was 281.7 (mg/m<sup>2</sup>/day). A test for the normal distribution of falling dust rates was conducted at each month of the study period (based on the

absence of statistically significant differences between the four stations) using the Shapiro- Wilk normality test. The results indicated that the distribution of data for all months of the study was normal. Therefore, one-way analysis of variance tests was conducted using Brown-Forsythe test and Tukey's multiple comparisons test. Test results showed no Existence Difference Statistically significant between the values of falling dust rates in the months, see figure 3 rates of dustfall in the months.



**Fig 3:** Dustfall rates at each month of the study period (based on the absence of statistically significant differences between the four stations)

It should be noted here that a comparison was made between the permissible limits and the values of the average rates, which was taken as the average of the months for each station of the study, and it was also taken as the average of the values of the dust rates for the four stations for each month. The result is that the values of falling dust rates in all months and at all stations that were studied exceeded the permissible values according to the British, Australian and Malaysian specifications, except for 4 stations, which were not observed in the same months. There was also an increase in falling dust rates above the permissible values according to American specifications in all months (except February) and in at least half of the stations studied. This indicates that there is one or more processes or activities that must be stopped until the dust levels are within the permissible limit.

**Discussion**

As can be seen from a table 2, the values of the rate of dustfall vary in months of the year, this may due to climatic factors such as relative humidity, rainfall, wind speed, and the topography of the region [18]. An increase in relative humidity leads to an increase in the size of dust particles, causing them to settle. In general, relative humidity levels vary depending on the change in outside temperature. Therefore, higher levels of relative humidity in summer than levels in winter may be a factor causing the values of falling dust rates in summer to be higher than those observed in winter in this study [19].

The dustfall standards permit industrial activity or any activity to add a certain amount of dust to the atmosphere, provided that the resulting atmospheric dustfall on the surfaces of the earth or properties does not exceed the values of the permissible parameters.

It should be noted here that a comparison was made between the permissible limits and the values of the average rates, which was taken as the average of the months for each station of the study, and it was also taken as the average of the values of the dust rates for the four stations for each month. The result is that the values of falling dust rates in all months and at all stations that were studied exceeded the permissible values according to the British, Australian and Malaysian specifications, except for 4 stations, which were not observed in the same months. There was also sand increase in falling dust rates above the permissible values according to American specifications in all months (except February) and in at least half of the stations studied. This indicates that there is one or more processes or activities that must be stopped until the dust levels are within the permissible limit.

As mentioned in the methodology section, the collected dust was mixed with distilled water then heated to remove soluble material. The findings indicate that most of the dust consists of insoluble material, which may include mineral content, such as soil dust. This mineral content may could be attributable to dust-producing activities such as cement, but could also be originated from other sources such as unsealed roads, etc.

**Conclusions and Recommendations**

There are operations, projects or activities that are responsible for producing dust and should not be allowed to produce a level of dust that exceeds the limits of rate of dustfall, including sources from industries, agriculture, unsealed roads, etc. The authorities must solve the problem of these operations or activities.

The study recommends that further studies be conducted to determine the physical properties and chemical composition

of baked fines, so that it is possible to estimate the potential dangers, whether to the environment, women, or property.

### Future works

After analysing the dustfall rate in four stations in Najaf, several areas for future research have been identified. Further study of the dust properties, such as color, to obtain important information about its origin is needed. Furthermore, identification of the chemical composition of the collected dust is required to assess the hazard.

### References

1. National Framework for Integrated Drought Risk Management in Iraq, Analytical Study [Internet]. c2014 Mar. Available from: [Insert URL if applicable]
2. Hazard Prevention and Control in the Work Environment: Airborne Dust [Internet]. [Insert date of publication]. Available from: [Insert URL if applicable]
3. Hazard Prevention and Control in the Work Environment: Airborne Dust [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
4. Dust Movement from and into Quarries [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
5. Air Quality & Haze Episodes in Malaysia [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
6. Heavy Metals in Dust Deposition in the Vicinity of Coal Ash Disposal Site Divkovic II [Internet]. [Insert date of publication]. Available from: [Insert URL if applicable]
7. Final Report on Dust Monitoring near Kellingley Coal Mine, North Yorkshire [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
8. Keller EA. Environmental Geology - 9<sup>th</sup> Edition. [Insert place of publication]; [insert publisher]; [insert year of publication].
9. Hazard Prevention and Control in the Work Environment: Airborne Dust [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
10. African Dust Contributions to Mean Ambient PM10 Mass-levels Across the Mediterranean Basin [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
11. Long-range Transport of Dust Aerosols over the Arabian Sea and Indian Region: A Case Study Using Satellite Data and Ground-based Measurements [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
12. Dustfall Associated with Dust Storms in the Al- Ahsa Oasis of Saudi Arabia [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
13. Dust Movement from and into Quarries [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
14. Dust Determination Methods and Instruments [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
15. Guide to Principles of Statistical Analysis Methodology and Quality Guides - Guide No. (10) [Internet]. ae.scad.www. [insert date of publication]. Available from: [insert URL if applicable]
16. Technical Appendix D – Air Quality and Dust [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
17. Correlation between Particulate Matter Concentrations and Meteorological Parameters at a Site in Ile-Ife, Nigeria [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
18. Evaluation of the Relationship between PM10 Concentrations and Heavy Metals during Normal and Dusty Days in Ahvaz, Iran [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]
19. Characterization and Elemental Composition of Atmospheric Aerosol Loads During Springtime Dust Storm in Western Saudi Arabia [Internet]. [Insert date of publication]. Available from: [insert URL if applicable]